Explosion speed of continuous state branching processes indexed by the Esscher transform

Loïc Chaumont* Clément Lamoureux*

March 19, 2025

Abstract

A branching process Z is said to be non conservative if it hits ∞ in a finite time with positive probability. It is well known that this happens if and only if the branching mechanism φ of Z satisfies $\int_{0+} d\lambda/|\varphi(\lambda)| < \infty$. We construct on the same probability space a family of conservative continuous state branching processes $Z^{(\varepsilon)}$, $\varepsilon \ge 0$, each process $Z^{(\varepsilon)}$ having $\varphi^{(\varepsilon)}(\lambda) = \varphi(\lambda + \varepsilon) - \varphi(\varepsilon)$ as branching mechanism, and such that the family $Z^{(\varepsilon)}$, $\varepsilon \ge 0$ converges a.s. to Z, as $\varepsilon \to 0$. Then we study the speed of convergence of $Z^{(\varepsilon)}$, when $\varepsilon \to 0$, referred to here as the explosion speed. More specifically, we characterize the functions f with $\lim_{\varepsilon \to 0} f(\varepsilon) = \infty$ and such that the first passage times $\sigma_{\varepsilon} = \inf\{t : Z_t^{(\varepsilon)} \ge f(\varepsilon)\}$ converge toward the explosion time of Z. Necessary and sufficient conditions are obtained for the weak convergence and convergence in L^1 . Then we give a sufficient condition for the almost sure convergence.

Keywords Continuous state branching process, spectrally positive Lévy process, Essher transform, Lamperti transform, first passage time

1 Introduction

Continuous state branching processes (CSBP) have two absorbing states: 0 and ∞ . We say that extinction (resp. explosion) occurs if the state 0 (resp. ∞) is attained in a finite time. Unlike the extinction properties which have been intensively studied in the past, the explosion of CSBP's does not seem to have been the subject of much research. The extinction and explosion conditions stated in Theorems 2.1 and 2.2 of the following section, however, reveal a sort of duality between these two phenomena. Our searches in the existing literature on the explosion of CSBP's have only led us to very few articles, most of them being quite recent: Li [10], Li and Zhou [9], Li, Foucart and Zhou [7] and some references therein.

Both articles [9] and [7] actually consider the more general case of nonlinear CSBP's. In the first one it is proved that when the process is non conservative (that is explosion occurs with positive probability) on its event of explosion, it tends to infinity in a finite time asymptotically along some deterministic curve. This speed of explosion is also characterised by the speed of convergence of the process of first passage times toward the explosion time. The second article provides sharper results in the case where the branching mechanism is regularly varying.

In the present article, we propose a quite different way of studying the explosion of a CSBP, Z, by constructing, on the same probability space, a sequence of conservative

^{*}Univ Angers, CNRS, LAREMA, SFR MATHSTIC, F-49000 Angers, France loic.chaumont@univ-angers.fr, clement.lamoureux@univ-angers.fr

processes $(Z^{(n)})$ that converges to the process Z. Then instead of considering the speed of explosion of the process Z itself, we study the speed of convergence of $(Z^{(n)})$ toward Z. This speed is characterised, on the set of explosion of Z, by the speed of convergence of the first passage times $\sigma_{v_n}^{(n)}$ of $Z^{(n)}$ above level v_n , for some sequence (v_n) that tends to ∞ . Intuitively, if (v_n) tends to ∞ not too fast, then $(\sigma_{v_n}^{(n)})$ is expected to converge to the explosion time ζ of Z whereas if it is too fast, then $(\sigma_{v_n}^{(n)})$ should tend to ∞ . We shall study the weak convergence, the convergence in L^1 norm and the almost sure convergence. Then we shall see that in the case of weak convergence an unexpected phenomenon occurs: when the speed of (v_n) belongs to some critical domain, the weak limit of $(\sigma_{v_n}^{(n)})$ is neither ζ nor ∞ but the law of $\zeta + c$, where the constant c > 0 can be considered as some 'residual mass'. This phenomenon is specific only to the weak convergence.

There are many ways of constructing a sequence of conservative branching processes $(Z^{(n)})$ that converges to some non conservative process Z, but perhaps the most natural way is to define $Z^{(n)}$ with branching mechanism $\varphi^{(n)}(\lambda) := \varphi(\varepsilon_n + \lambda) - \varphi(\varepsilon_n)$, where $\varepsilon_n \downarrow 0$ and φ is the branching mechanism of Z. Then the weak convergence of $(Z^{(n)})$ toward Z clearly holds. Moreover, it is possible to construct the sequence $(Z^{(n)})$ on the same probability space as Z. This coupling is one of the main results of the present paper. We obtain the sequence $(Z^{(n)})$ by first constructing an increasing sequence of Lévy processes $(X^{(n)})$ such that each process $X^{(n)}$ has branching mechanism $\varphi^{(n)}$ defined above, so that the law of $X^{(n)}$ is an Esscher transform of the law of X. Then $Z^{(n)}$ is obtained by the well known means of the Lamperti transformation. This construction ensures an a.s. uniform convergence of $X^{(n)}$ toward X and an a.s. convergence of $Z^{(n)}$ toward Z in the Skohorod's topology.

The next section is devoted to some basic notions on branching processes as well as the construction of the sequence $(Z^{(n)})$. Then we state our main results on the explosion speed of $(Z^{(n)})$ in Section 3 and these results are proved in Section 4.

2 A family of CSBP indexed by the Esscher transform

2.1 A brief review of CSBP's

A continuous state branching process (CSBP) $(Z_t, t \ge 0)$ is a $[0, \infty]$ -valued Markov process with probabilities (P_x) satisfying the following property, called the branching property,

$$E_{x_1+x_2}(e^{-\lambda Z_t}) = E_{x_1}(e^{-\lambda Z_t})E_{x_2}(e^{-\lambda Z_t}), \quad x_1, x_2, \lambda, t \ge 0.$$

This property implies that the states 0 and ∞ are absorbing, see [5]. Moreover there exists a differentiable function $u_t : [0, \infty) \to [0, \infty)$, called the Laplace exponent of Z, which satisfies

$$\mathbb{E}_x(e^{-\lambda Z_t}) = e^{-xu_t(\lambda)}, \quad x, \lambda, t \ge 0.$$
(2.1)

Then the Markov property yields the following semi-group property of u_t ,

$$u_{t+s}(\lambda) = u_t(u_s(\lambda)), \quad s, t, \lambda \ge 0,$$

from which we derive that u_t solves the differential equation,

$$\frac{\partial u_t}{\partial t}(\lambda) + \varphi(u_t(\lambda)) = 0 \quad ; \quad u_0(\lambda) = \lambda \,, \tag{2.2}$$

where $\varphi : [0, \infty) \to \mathbb{R}$ is the Laplace exponent of a spectrally positive Lévy process (spLp) that we will denote $(X_t, t \ge 0)$, that is

$$\mathbb{E}(e^{-\lambda X_t}) = e^{t\varphi(\lambda)}, \quad \lambda \ge 0.$$

We denote by $\mathbb{P}_x, x \in \mathbb{R}$ a family of probability measures under which X starts from x and we set $\mathbb{P}_0 := \mathbb{P}$. The function φ is called the branching mechanism of Z. It is a log-convex function such that $\lim_{\lambda \to \infty} \varphi(\lambda) = \infty$ if X is not a subordinator and $\varphi \leq 0$ when X is a subordinator. According to the Lévy-Khintchine formula, φ can be expressed as,

$$\varphi(\lambda) = -q + a\lambda + \frac{1}{2}\sigma^2\lambda^2 + \int_{(0,\infty)} \left(e^{-\lambda x} - 1 + \lambda x \mathbb{1}_{\{x<1\}}\right) \pi(dx), \qquad (2.3)$$

where $q \ge 0$, $a \in \mathbb{R}$, $\sigma \ge 0$ and π is a Lévy measure, that is, since X has no positive jumps, a measure on $(0, \infty)$ such that $\int_{(0,\infty)} (x^2 \wedge 1)\pi(dx) < \infty$. We can check that for all $\lambda > 0$, u_t is the unique solution of (2.2) and it is given for all $t \ge 0$ by,

$$\int_{u_t(\lambda)}^{\lambda} \frac{du}{\varphi(u)} = t.$$
(2.4)

Let us now focus on the asymptotic behaviour of CSBP's. We first define the following (disjoint) sets,

$$A_0 = \left\{ \lim_{t \to \infty} Z_t = 0 \right\}$$
 and $A_\infty = \left\{ \lim_{t \to \infty} Z_t = \infty \right\}$.

Note that each of the absorbing states 0 and ∞ can be attained by Z in a finite or infinite time. Define the largest root of the branching mechanism φ as,

$$\rho = \sup\{\lambda \ge 0 : \varphi(\lambda) \le 0\},\$$

and note that $\rho = \infty$ if and only if φ is the Laplace exponent of a subordinator. Then the events A_0 and A_∞ satisfy the dichotomy described in the two following theorems, see [5] and Chapter 12 of [6].

Theorem 2.1. Let Z be a CSBP with branching mechanism φ . Then $\mathbb{P}_x(A_0 \cup A_\infty) = 1$, for all $x \ge 0$. Moreover,

$$P_x(A_0) = e^{-x\rho}$$
 and $P_x(A_\infty) = 1 - e^{-x\rho}$.

In turn, each of the events A_0 and A_∞ can be partitioned in the following way: the event A_0 can be written as the union of two disjoint sets, $A_0 = A_0^{\rightarrow} \cup A_0^{\downarrow}$, where

$$A_0^{\rightarrow} = \left\{ \lim_{t \to \infty} Z_t = 0 \text{ and } Z_t > 0 \text{ for all } t > 0 \right\} \text{ and } A_0^{\downarrow} = \{ Z_t = 0, \text{ for some } t > 0 \} ,$$

and the event A_{∞} can be written as the union of two disjoint sets, $A_{\infty} = A_{\infty}^{\rightarrow} \cup A_{\infty}^{\uparrow}$, where

$$A_{\infty}^{\rightarrow} = \left\{ \lim_{t \to \infty} Z_t = \infty \text{ and } Z_t < \infty \text{ for all } t > 0 \right\} \text{ and } A_{\infty}^{\uparrow} = \left\{ Z_t = \infty, \text{ for some } t > 0 \right\}.$$

Then the following dichotomy holds for each of the events A_0 and A_{∞} .

Theorem 2.2. Let Z be a CSBP with branching mechanism φ . Then the following assertions regarding the event of extinction A_0^{\downarrow} are equivalent,

- (i) There is $\theta > 0$, such that $\int_{\theta}^{\infty} \frac{du}{|\varphi(u)|} < \infty$,
- (*ii*) for all $x \ge 0$, $P_x(A_0^{\downarrow}) = P_x(A_0) = e^{-x\rho}$ and $\rho < \infty$,
- (iii) for all t > 0, $u_t(\infty) < \infty$.

Similarly, for the event of explosion A^{\uparrow}_{∞} , the following assertions are equivalent,

- (j) There is $\theta > 0$, such that $\int_0^\theta \frac{du}{|\varphi(u)|} < \infty$,
- (*jj*) for all $x \ge 0$, $P_x(A_{\infty}^{\uparrow}) = P_x(A_{\infty}) = 1 e^{-x\rho}$ and $\rho > 0$,
- (jjj) for all t > 0, $u_t(0) > 0$.

2.2 Path construction of a family of CSBP's

Let us first recall the Esscher transform of a spLp. This can simply be expressed from the Laplace exponent φ of the process as follows: for all $\varepsilon \geq 0$, the function

$$\varphi^{(\varepsilon)}(\lambda) := \varphi(\lambda + \varepsilon) - \varphi(\varepsilon), \quad \lambda \ge 0,$$

remains the Laplace exponent of a spLp. In terms of martingale change of measure, the Esscher transform of a spLp X with Laplace exponent φ is (the law of) a spLp $X^{(\varepsilon)}$ with Laplace exponent $\varphi^{(\varepsilon)}$ given by

$$\mathbb{E}_x(F(X_s^{(\varepsilon)}, s \in [0, t])) = \mathbb{E}_x\left(F(X_s, s \in [0, t])e^{-\varepsilon X_t - t\varphi(\varepsilon)}\right)$$

for all t > 0 and all bounded, measurable functional F, see for instance Theorem 3.9 in [6].

Given the Laplace exponent φ of a spLp, through our next result, we show how to construct on the same probability space, an increasing family of spLp's $\{X^{(\varepsilon)}, \varepsilon \geq 0\}$ such that for each $\varepsilon \geq 0$, $X^{(\varepsilon)}$ has Laplace exponent $\varphi^{(\varepsilon)}$.

Theorem 2.3. Let φ be given by (2.3) and satisfying $\varphi(0) = 0$. Let $(\varepsilon_n)_{n\geq 0}$ be a decreasing sequence of real numbers such that $\lim_n \varepsilon_n = 0$.

Then on some probability space we can define an increasing sequence of Lévy processes $(X^{(\varepsilon_n)})_{n\geq 0}$, each process $X^{(\varepsilon_n)}$ having $\varphi^{(\varepsilon_n)}$ as Laplace exponent, and such that $(S^{(\varepsilon_n)})_{n\geq 0} := (X^{(\varepsilon_{n+1})} - X^{(\varepsilon_n)})_{n\geq 0}$ is a sequence of independent subordinators that is itself independent of $X^{(\varepsilon_0)}$. For each n, $S^{(\varepsilon_n)}$ has Laplace exponent,

$$\Gamma^{(\varepsilon_n)}(\lambda) = (\varepsilon_{n+1} - \varepsilon_n)\sigma^2\lambda + \int_{(0,\infty)} \left(e^{-\lambda x} - 1\right) \left(e^{-\varepsilon_{n+1}x} - e^{-\varepsilon_n x}\right)\pi(dx).$$
(2.5)

Moreover the sequence $(X^{(\varepsilon_n)})$ converges uniformly over any closed intervals of \mathbb{R}_+ , almost surely, toward a Lévy process X whose Laplace exponent is φ .

Let us now recall the Lamperti representation of a CSBP with branching mechanism φ from the path of a spectrally positive Lévy process X with Laplace exponent φ . This is the mapping defined as follows:

$$L(X)_t := \begin{cases} X(I_t \wedge \tau), & \text{if } I_t \wedge \tau < \infty, \\ \infty, & \text{if } I_t \wedge \tau = \infty, \end{cases}$$
(2.6)

where $I_t = \inf \left\{ s : \int_0^s \frac{du}{X_u} > t \right\}$ and $\tau = \inf \{ t \ge 0 : X_t \le 0 \}$. Note that in (2.6), for t > 0and $X_0 > 0$, both variables I_t and τ can be infinite only if X drifts to ∞ . Then under \mathbb{P}_x , $x \ge 0$, the process $(L(X)_t, t \ge 0)$ is a CSBP with branching mechanism φ , issued from x, see for instance Theorem 12.2, p.337 in [6]. Note also that the transformation (2.6) is invertible and the paths of the process $(X_t, 0 \le t \le \tau)$ can be recovered from those of the process Z.

Let us fix a Laplace exponent φ as in (2.3) and a decreasing sequence (ε_n) of real numbers such that $\lim_n \varepsilon_n = 0$. Then on some probability space, we define Lévy processes X and $X^{(\varepsilon_n)}$, $n \ge 0$ as in Proposition 2.3 and to make our notations simpler, we set,

$$\varphi^{(n)} := \varphi^{(\varepsilon_n)}$$
 and $X^{(n)} := X^{(\varepsilon_n)}$

Furthermore from the paths of the sequence $(X^{(n)}, n \ge 0)$ and those of its limit X, we define the CSBP's $(Z^{(n)}, n \ge 0)$ and Z through the Lamperti transformation:

$$Z^{(n)} := L(X^{(n)}), \ n \ge 0 \text{ and } Z := L(X).$$
 (2.7)

Given the convergence result obtained in Theorem 2.3, it is natural to wonder if the sequence $(Z^{(n)}, n \ge 0)$ converges toward Z in some sense. This is the purpose of the next theorem which follows directly from Corollary 6 in [3]. We shall denote by ζ the first hitting time of ∞ by the process Z, that is,

$$\zeta = \inf\{t \ge 0 : Z_t = \infty\},\tag{2.8}$$

where $\inf \emptyset = \infty$ (according to our notation $\{\zeta < \infty\} = A_{\infty}^{\uparrow}$). Note that under \mathbb{P}_x , the processes $Z^{(n)}$, $n \ge 0$ and Z are issued from x as well as X and $X^{(n)}$, $n \ge 0$. Then from now on, we will only use the probabilities \mathbb{P}_x , $x \ge 0$ for the processes X, $X^{(n)}$, Z and $Z^{(n)}$.

Theorem 2.4. For all t, x > 0, the family of processes $(Z_s^{(n)}, 0 \le s \le t)$ converges \mathbb{P}_x a.s. on the set $\{t < \zeta\}$ in the Skohorod's J_1 topology toward the process $(Z_s, 0 \le s \le t)$.

3 On the speed of explosion of the sequence $\{Z^{(n)}, n \ge 0\}$

3.1 Presentation of the problem

Throughout the remainder of this article, we will assume that φ is the Laplace exponent of a spLp satisfying the condition,

$$\varphi(0) = 0$$
 and there is $\theta > 0$, such that $\int_0^\theta \frac{du}{|\varphi(u)|} < \infty.$ (3.1)

Then like in the previous section, we fix a decreasing sequence (ε_n) such that $\lim_{n\to\infty} \varepsilon_n = 0$ and we construct an increasing sequence of Lévy processes $(X^{(n)})$ with respective Laplace exponents $\varphi^{(n)}(\lambda) := \varphi(\varepsilon_n + \lambda) - \varphi(\varepsilon_n), \lambda \ge 0$, as in Theorem 2.3. Its limit, X, is then a Lévy process with Laplace exponent φ . Note that all the processes $X^{(n)}$, $n \ge 0$ and X drift to ∞ . Recall also the construction (2.7), from the paths of $X^{(n)}$, $n \ge 0$ and X, of the branching processes $Z^{(n)}$, $n \ge 0$ and their limit Z whose respective branching mechanism are $\varphi^{(n)}$, $n \ge 0$ and φ . We emphasize that, due to the time change in the transformation (2.7), the sequence $(Z^{(n)})$ is no longer monotone.

From Theorem 2.2 and our assumption (3.1), explosion occurs for the process Z, that is:

for all
$$x > 0$$
, $\mathbb{P}_x(\zeta < \infty) = \mathbb{P}_x(A_\infty^{\uparrow}) = \mathbb{P}_x(A_\infty) = 1 - e^{-x\rho} > 0$

where ζ is the explosion time defined in (2.8). Moreover, it can be seen from the Lamperti transformation (2.6) that Z explodes in a 'continuous way', that is $Z_{\zeta-} = \infty$, \mathbb{P}_x -a.s. for all x > 0 on the set $\{\zeta < \infty\}$, see Figure 1 below. Let us also point out on the fact that,

from Theorem 2.3, the sequence of sets $\{\tau^{(n)} = \infty\}$ is increasing and tends to $\{\tau = \infty\}$, where $\tau^{(n)} = \inf\{t \ge 0 : X_t^{(n)} \le 0\}$ and $\tau = \inf\{t \ge 0 : X_t \le 0\}$. Moreover it appears from the construction (2.7) that

$$\{\zeta < \infty\} = \{\tau = \infty\}.$$

On the other hand, since for all $n \ge 0$, $|\varphi^{(n)'}(0)| = |\varphi'(\varepsilon_n)| < \infty$, Theorem 2.2 implies that explosion occurs for none of the processes $Z^{(n)}$, that is with obvious notation for all $n \ge 0$ and $x \ge 0$, $\mathbb{P}_x(A_{\infty}^{n,\uparrow}) = 0$. Note also that from (3.1), $\varphi'(0) = -\infty$, so that we can assume without loss of generality that $\varphi'(\lambda) < 0$, for all $\lambda \in [0, \varepsilon_0]$. This implies that for all $n \ge 0$, $\varphi^{(n)'}(0) = \varphi'(\varepsilon_n) < 0$, so that $\rho_n := \sup\{\lambda \ge 0 : \varphi^{(n)}(\lambda) \le 0\} > 0$. Therefore, each process $Z^{(n)}$ tends to ∞ with positive probability. This can be summarized as follows:

for all $x \ge 0$ and $n \ge 0$, $\mathbb{P}_x(A_{\infty}^{n,\uparrow}) = 0$ and $\mathbb{P}_x(A_{\infty}^n) = 1 - e^{-x\rho_n} > 0$.

Let us now denote by $\sigma_y^{(n)}$ the first passage time by the process $Z^{(n)}$ above a level y > 0, that is

$$\sigma_y^{(n)} = \inf\{t : Z_t^{(n)} \ge y\},\$$

and recall that from Theorem 2.4, for all t, x > 0, the sequence of non explosive processes $(Z_s^{(n)}, 0 \le s \le t), n \ge 0$, converges \mathbb{P}_x -a.s. on the set $\{t < \zeta\}$ toward the explosive process $(Z_s, 0 \le s \le t)$ in the Skohorod's J_1 topology. In order to study the speed of explosion of the family $(Z^{(n)}, n \ge 0)$, we shall characterise the set of increasing sequences (v_n) such that $\lim_{n\to\infty} v_n = \infty$ and for which the sequence of random variables $(\sigma_{v_n}^{(n)})$ converges in some sense. If (v_n) is not too fast, then $(\sigma_{v_n}^{(n)})$ is expected to converge, whereas if the speed of convergence of (v_n) is too fast compared to this of (ε_n) , then $\sigma_{v_n}^{(n)} \to \infty$ should go to ∞ . Therefore, there must be a threshold for this speed under which $\sigma_{v_n}^{(n)}$ tends toward a non degenerate random variable which is actually expected to be the explosion time ζ of Z, see Figure 1 below.



Figure 1: Z in red, $Z^{(n)}$ in orange and $Z^{(m)}$ in blue for n > m.

3.2 Convergence in distribution

In the remainder of this paper, we shall often use the notation,

$$\phi := -\varphi$$
 and $\phi^{(n)} := -\varphi^{(n)}$

so that ϕ and $\phi^{(n)}$ are positive, concave functions on $(0, \rho)$ and $(0, \rho_n)$, respectively, where $\rho_n = \sup\{\lambda \ge 0 : \varphi^{(n)} \le 0\}$. Moreover the sequence of functions $(\phi^{(n)})$ is increasing and satisfies $\lim_{n\to\infty} \phi^{(n)} = \phi$. We denote the Laplace exponent of each process $Z^{(n)}$ by $u^{(n)}$, see (2.1). In all the following statements, the function $h: \mathbb{Z}_+ \to \mathbb{R}_+$ will always satisfy,

$$(h(n))_{n\geq 0}$$
 is a decreasing sequence such that $\lim_{n\to\infty} h(n) = 0.$ (3.2)

Let us also specify that when mentioning the existence of the limit of a non negative valued sequence, we mean that it exists in $[0, \infty]$.

Theorem 3.1. The following assertions are equivalent:

1. There is a $[0, \infty]$ -valued random variable Y such that for all k > 0 and for all $x \ge 0$, under \mathbb{P}_x , $\sigma_{k/h(n)}^{(n)} \xrightarrow[n \to \infty]{(d)} Y$.

2. For all
$$\theta \in (0, \rho)$$
, $\lim_{n \to \infty} \int_{h(n)}^{\theta} \frac{du}{\phi^{(n)}(u)}$ exists.

3. For all $t \ge 0$, $\lim_{n\to\infty} u_t^{(n)}(h(n)) := l_t(h)$ exists.

When the above conditions are satisfied, the mapping $t \mapsto l_t(h)$ is continuous and non decreasing on $[0, \infty)$. It satisfies $l_0(h) = 0$ and takes its values in $[0, \rho)$. Moreover, for all $x \ge 0$, under \mathbb{P}_x ,

$$Y \stackrel{\scriptscriptstyle (d)}{=} \zeta + c(h) \quad and \quad \lim_{n \to \infty} \int_{h(n)}^{\theta} \frac{du}{\phi^{(n)}(u)} = \int_{0}^{\theta} \frac{du}{\phi(u)} + c(h) \,,$$

where the constant c(h) is given by $c(h) = \sup\{t \ge 0 : l_t(h) = 0\}$.

Note that under our assumptions $\mathbb{P}_x(\zeta < \infty) > 0$, hence the random variable Y involved in Theorem 3.1 satisfies $\mathbb{P}_x(Y = \infty) = 1$ if and only if $c(h) = \infty$. The following remark can be understood as a complement to Theorem 3.1.

Remark 3.1. Given a constant $c \in [0, \infty]$ and $\theta \in (0, \rho)$, one can always construct a decreasing function h_c such that

$$\lim_{n \to \infty} \int_{h_c(n)}^{\theta} \frac{du}{\phi^{(n)}(u)} = \int_0^{\theta} \frac{du}{\phi(u)} + c.$$
(3.3)

To this end, note that the mapping $x \mapsto F_n(x) := \int_x^\theta \frac{du}{\phi^{(n)}(u)}$ defines a decreasing bijection from $(0,\theta)$ to $(0,\infty)$. Let us denote by F_n^{-1} its inverse. Then for $c \in [0,\infty)$, the sequence $h_c(n) := F_n^{-1} \left(\int_0^\theta \frac{du}{\phi(u)} + c \right)$ fulfils condition (3.3) and for $c = \infty$, we can take for instance $h_c(n) := F_n^{-1} \left(\int_0^\theta \frac{du}{\phi(u)} + n \right).$ Theorem 3.1 induces the following classification of sequences:

$$\begin{aligned} \mathcal{Z} &:= \{h : (h(n))_{n \ge 0} \text{ satisfies conditions } 1., 2. \text{ and } 3. \text{ of Theorem } 3.1\} \\ \mathcal{Z}_0 &:= \{h \in \mathcal{Z} : c(h) = 0\}, \quad \mathcal{Z}_\infty := \{h \in \mathcal{Z} : c(h) = \infty\} \\ \mathcal{Z}_c &:= \{h \in \mathcal{Z} : 0 < c(h) < \infty\}. \end{aligned}$$

We shall prove in Lemma 4.7 below that a sufficient condition for $h \in \mathbb{Z}_0$ is,

$$\liminf_{n \to \infty} \phi(h(n) + \varepsilon_n) / \phi(\varepsilon_n) > 1.$$
(3.4)

Except in the regularly varying case, see Theorem 3.2 below, it is not clear that the weak convergence of $(\sigma_{k/h(n)}^{(n)})$ for some k > 0 implies this convergence for all k > 0, that is $h \in \mathbb{Z}_0$. However, if this convergence holds when k is replaced by an independent exponentially distributed random variable, then $h \in \mathbb{Z}_0$ as shown in the following result.

Proposition 3.1. Let e be an exponentially distributed random variable with parameter 1 which is independent of the sequence $(Z^{(n)}, n \ge 1)$. Then $\sigma_{e/h(n)}^{(n)} \xrightarrow{(d)} \zeta$, if and only if $h \in \mathbb{Z}_0$.

The next results allow us to compare the limits of $(\sigma_{1/h(n)}^{(n)})$ and $(\sigma_{1/\tilde{h}(n)}^{(n)})$ according to the relative behaviours of the sequences (h(n)) and $(\tilde{h}(n))$.

Proposition 3.2. Let $\tilde{h} : \mathbb{Z}_+ \to \mathbb{R}_+$ be such that $(\tilde{h}(n))$ is decreasing with $\lim_{n\to\infty} \tilde{h}(n) = 0$. If $\sigma_{1/h(n)}^{(n)} \xrightarrow{(d)} \zeta$ and $h(n) \leq \tilde{h}(n)$, for all n sufficiently large, then $\sigma_{1/\tilde{h}(n)}^{(n)} \xrightarrow{(d)} \zeta$.

Proposition 3.3. Let $h \in \mathbb{Z}$ and $\tilde{h} : \mathbb{Z}_+ \to \mathbb{R}_+$ be such that $(\tilde{h}(n))$ is a decreasing sequence with $\lim_{n\to\infty} \tilde{h}(n) = 0$.

- 1. If $\tilde{h} \simeq h$, then $\tilde{h} \in \mathcal{Z}$ and $c(\tilde{h}) = c(h)$.
- 2. If $h \in \mathcal{Z}_c$ and if $\tilde{h} \in \mathcal{Z}$ satisfies $h(n) \leq \tilde{h}(n)$ for all n sufficiently large, then $c(\tilde{h}) \leq c(h)$.
- 3. If $h \in \mathbb{Z}_0$ and if \tilde{h} satisfies $h(n) \leq \tilde{h}(n)$ for all n sufficiently large, then $\tilde{h} \in \mathbb{Z}_0$.
- 4. If $h \in \mathbb{Z}_{\infty}$ and if \tilde{h} satisfies $\tilde{h}(n) \leq h(n)$ for all n sufficiently large, then $\tilde{h} \in \mathbb{Z}_{\infty}$.

We can see in assertion 2. of Proposition 3.3 that if $h \in \mathbb{Z}_c$, then it is not enough that $h \leq \tilde{h}$ or $\tilde{h} \leq h$ to deduce something on \tilde{h} , contrary to what is asserted in parts 3. and 4. Actually the proof of assertion 2. requires the additional hypothesis that $\tilde{h} \in \mathbb{Z}$.

The following proposition suggests that the existence of the limit of $|\log h(n)|/\phi'(\varepsilon_n)$ as n tends to ∞ might be a better criterion for the weak convergence of $(\sigma_{k/h(n)}^{(n)})$ than the convergence of $\int_{h(n)}^{\theta} \frac{du}{\phi^{(n)}(u)}$, as stated in Theorem 3.1. This can actually be proved in the regularly varying case, see Theorem 3.2.

Proposition 3.4.

1. If $\lim_{n \to \infty} \frac{|\log h(n)|}{\phi'(\varepsilon_n)} = \infty$, then $h \in \mathbb{Z}_{\infty}$. 2. If $h \in \mathbb{Z}_0$, then $\lim_{n \to \infty} \frac{|\log h(n)|}{\phi'(\varepsilon_n)} = 0$. In the case where φ is regularly varying at 0, a better criterion allows us to characterize the limit of $\sigma_{1/h(n)}^{(n)}$.

Theorem 3.2. Suppose that φ is regularly varying at 0. Then the following assertions are equivalent:

1. The limit
$$c(h) := \lim_{n \to \infty} \frac{|\log h(n)|}{\phi'(\varepsilon_n)}$$
 exists.

2. For all
$$k > 0$$
, $\sigma_{k/h(n)}^{(n)} \xrightarrow{(a)} \zeta + c(h)$, that is $h \in \mathbb{Z}$.

When these assumptions are satisfied, the constant the constant c(h) is the same as that in Theorem 3.1. Moreover assertions 1. and 2. for c(h) = 0 are equivalent to the following one:

3. For some
$$k > 0$$
, $\sigma_{k/h(n)}^{(n)} \xrightarrow[n \to \infty]{} \zeta$.

3.3 Convergence in L_1 and almost sure convergence

We recall the usual convention $0 \times \infty = \infty \times 0 = 0$, so that in particular, if Y is any non negative random variable, then $Y \mathbb{1}_{\{Y < \infty\}}$ is a.s. finite. As we will see in this section, when $h \in \mathbb{Z}_0$, stronger convergences than weak convergence actually hold. Let us first notice that the time ζ admits all its moments on the set $\{\zeta < \infty\}$. This result is also proved in Theorem 3.1 of [9] where the moments of ζ are expressed in a different form, see also Theorems 1.5 and 1.6 in [10] where this result is obtained in a particular case.

Proposition 3.5. For all $n \in \mathbb{Z}_+$ and x > 0, $\mathbb{E}_x\left(\zeta^n \mathbb{1}_{\{\zeta < \infty\}}\right) < \infty$. Moreover for all $n \in \mathbb{Z}_+$, x > 0 and $\lambda \ge 0$,

$$\mathbb{E}_x\left(\zeta^n \mathbb{1}_{\{\zeta<\infty\}}\right) = x \int_0^\rho F(y)^n e^{-xy} dy \quad and \quad \mathbb{E}_x\left(e^{-\lambda\zeta}\right) = x \int_0^\rho e^{-\lambda F(y)} e^{-xy} dy,$$

where $F(y) = \int_0^y \frac{du}{\phi(u)}$, for $y \in [0, \rho)$. In particular, $\mathbb{E}_x\left(\zeta \mathbb{1}_{\{\zeta<\infty\}}\right) = \int_0^\rho \frac{e^{-\lambda x} - e^{-\rho x}}{\phi(\lambda)} d\lambda.$

As we will show in Subsection 4.3.1, the first passage times $\sigma_{1/h(n)}^{(n)}$ are integrable on the

set $\{\sigma_{1/h(n)}^{(n)} < \infty\}$ and we will prove the following result.

Theorem 3.3. Assume that $h \in \mathbb{Z}_0$. Then for all x > 0 and $n \ge 0$, the random variable $\sigma_{1/h(n)}^{(n)} \mathbb{1}_{\{\sigma_{1/h(n)}^{(n)} < \infty\}}$ is integrable under \mathbb{P}_x , moreover,

$$\sigma_{1/h(n)}^{(n)} \mathbb{1}_{\{\sigma_{1/h(n)}^{(n)} < \infty\}} \xrightarrow{\mathrm{L}^1(\mathbb{P}_x)}{n \to \infty} \zeta \mathbb{1}_{\{\zeta < \infty\}}.$$

It follows from Theorems 3.1 and 3.3 that for $h \in \mathbb{Z}_0$, the convergence in law of $\left(\sigma_{1/h(n)}^{(n)}\right)$ toward ζ is equivalent to the convergence in L^1 of $\left(\sigma_{1/h(n)}^{(n)}\mathbb{1}_{\{\sigma_{1/h(n)}^{(n)}<\infty\}}\right)$ toward $\zeta\mathbb{1}_{\{\zeta<\infty\}}$.

We also emphasize that if $h \in \mathbb{Z}_c$, that is if $\sigma_{1/h(n)}^{(n)} \xrightarrow[n \to \infty]{(d)} \zeta + c(h)$, with $c(h) \in (0, \infty)$, then the convergence in L^1 of $\left(\sigma_{1/h(n)}^{(n)} \mathbb{1}_{\{\sigma_{1/h(n)}^{(n)} < \infty\}}\right)$ toward $\zeta \mathbb{1}_{\{\zeta < \infty\}} + c(h)$ would make sense only if the constant c(h) can be obtained as a functional of the paths of the sequence $(Z^{(n)})$ and those of the process Z. This does not seem very realistic and the constant c(h) may just be an adjustment value that appears in the weak limit, when h belongs to the critical domain \mathcal{Z}_c . The same remark obviously holds for almost sure convergence, below.

We now state that under a stronger assumption than this of Theorem 3.3, the sequence $\left(\sigma_{1/h(n)}^{(n)} \mathbb{I}_{\{\sigma_{1/h(n)}^{(n)} < \infty\}}\right)$ converges almost surely toward $\zeta \mathbb{I}_{\{\zeta < \infty\}}$.

Theorem 3.4. Assume that $\sum \phi(\varepsilon_n)/\phi(\varepsilon_n + h(n)) < \infty$. Then for all x > 0,

$$\sigma_{1/h(n)}^{(n)} \mathbb{1}_{\{\sigma_{1/h(n)}^{(n)} < \infty\}} \xrightarrow{\mathbb{P}_x - a.s.}{n \to \infty} \zeta \mathbb{1}_{\{\zeta < \infty\}}.$$

The condition $\sum \phi(\varepsilon_n)/\phi(\varepsilon_n + h(n)) < \infty$ is not a necessary condition for the almost sure convergence of $\binom{n}{1/h(n)}$ to hold. Indeed, take for instance $\phi(\lambda) = k\lambda^{1/2}, \lambda \ge 0$, where k is some positive constant. Take also $h(n) = \varepsilon_n = n^{-1}$. Then from Theorems 3.2 and 3.3, there is a sequence of integers $(n_i)_{i\ge 0}$ with $n_i \to \infty$, as $i \to \infty$ such that $\sigma_{1/h(n_i)}^{(n_i)}$ converges almost surely toward ζ , as $i \to \infty$. However $\sum_i \phi(\varepsilon_{n_i})/\phi(\varepsilon_{n_i} + h(n_i)) = \infty$. This provides a counterexample with $\phi(\lambda) = k\lambda^{1/2}, \tilde{\varepsilon}_i := \varepsilon_{n_i}$ and $\tilde{h}(i) := h(n_i)$.

From Theorem 3.4, convergence in distribution of $(\sigma_{1/h(n)}^{(n)})$ holds, that is $h \in \mathbb{Z}_0$, under the condition $\sum \phi(\varepsilon_n)/\phi(\varepsilon_n + h(n)) < \infty$. This can be checked directly by applying Lemma 4.7, see (3.4) above.

We end this section with a related result that provides another way to evaluate the speed of convergence of the sequence $(Z^{(n)})$ toward Z. Let us give ourselves an exponentially distributed random variable e with parameter 1 that is independent of the sequence of Lévy processes $(X^{(n)})$. For each $n \ge 0$, denote by $\tilde{X}^{(n)}$ the Lévy process $X^{(n)}$ hilled at the independent exponential time $e_n := e/h(n)$. The killed Lévy process $\tilde{X}^{(n)}$ has Laplace exponent $\tilde{\varphi}^{(n)}(\lambda) := \varphi^{(n)}(\lambda) - h(n)$. Then we define the sequence $\tilde{Z}^{(n)} := L(\tilde{X}^{(n)}), n \ge 0$ of branching processes obtained from the sequence of Lévy processes $\tilde{X}^{(n)}$ through the Lamperti representation (2.6). The process $\tilde{Z}^{(n)}$ has branching mechanism $\tilde{\varphi}^{(n)}$. Recall from Theorem 2.2 that $\tilde{Z}^{(n)}$ hits ∞ in a finite time with positive probability and from (2.6) this is done through a jump, that is $\tilde{Z}^{(n)}_{\tilde{\zeta}^{(n)}} < \infty$, \mathbb{P}_x -a.s. on the set $\{\tilde{\zeta}^{(n)} < \infty\}$, for all x > 0, where $\tilde{\zeta}^{(n)} := \inf \left\{ t : \tilde{Z}^{(n)}_t = \infty \right\}$. Actually, the process $\tilde{Z}^{(n)}$ corresponds to the process $Z^{(n)}$ killed at the time $\int_0^{e_n \wedge \tau^{(n)}} \frac{du}{X^{(n)}_u}$, where $\tau^{(n)} := \inf \left\{ t : X^{(n)}_t \le 0 \right\}$. Moreover the hitting time $\tilde{\zeta}^{(n)}$ satisfies

$$\tilde{\zeta}^{(n)} \mathbb{1}_{\{\tilde{\zeta}^{(n)} < \infty\}} = \int_0^{\mathbf{e}_n} \frac{du}{X_u^{(n)}} \mathbb{1}_{\{\mathbf{e}_n < \tau^{(n)}\}}.$$
(3.5)

By reinforcing the hypothesis of Theorem 3.4, we can prove that the sequence of explosion times $(\tilde{\zeta}^{(n)} \mathbb{1}_{\{\tilde{\zeta}^{(n)} < \infty\}})$ converges almost surely toward $\zeta \mathbb{1}_{\{\zeta < \infty\}}$.

Proposition 3.6. If $\sum \phi(\varepsilon_n)/h(n) < \infty$, then for all x > 0,

$$\tilde{\zeta}^{(n)} 1\!\!1_{\{\tilde{\zeta}^{(n)} < \infty\}} \xrightarrow{\mathbb{P}_x - a.s.}{n \to \infty} \zeta 1\!\!1_{\{\zeta < \infty\}}$$

Proposition 3.6 raises the question of the convergence in L^1 of the sequence $(\tilde{\zeta}^{(n)} \mathbb{1}_{\{\tilde{\zeta}^{(n)} < \infty\}})$. When X is a subordinator and under the sole assumption that $h \in \mathbb{Z}_0$, this is a direct consequence of Corollary 4.1. The general case seems much more delicate.

4 Proofs

4.1 Proof of Theorem 2.3

Let us first write $\varphi^{(\varepsilon_n)}$ as follows,

$$\begin{split} \varphi^{(\varepsilon_n)}(\lambda) &= \varphi(\lambda + \varepsilon_n) - \varphi(\varepsilon_n) \\ &= \left(a + \varepsilon_n \sigma^2 + \int_{(0,\infty)} x \left(1 - e^{-\varepsilon_n x} \right) \mathbb{1}_{\{x < 1\}} \pi(dx) \right) \lambda + \frac{1}{2} \sigma^2 \lambda^2 \\ &+ \int_{(0,\infty)} \left(e^{-\lambda x} - 1 + \lambda x \mathbb{1}_{\{x < 1\}} \right) e^{-\varepsilon_n x} \pi(dx) \,. \end{split}$$

This yields that the difference

$$\Gamma^{(\varepsilon_n)}(\lambda) := \varphi^{(\varepsilon_{n+1})}(\lambda) - \varphi^{(\varepsilon_n)}(\lambda)$$

= $(\varepsilon_{n+1} - \varepsilon_n)\sigma^2\lambda + \int_{(0,\infty)} \left(e^{-\lambda x} - 1\right) \left(e^{-\varepsilon_{n+1}x} - e^{-\varepsilon_nx}\right)\pi(dx)$

is the Laplace exponent of a subordinator.

Then on the same probability space, define a Lévy process $X^{(\varepsilon_0)}$ with Laplace exponent $\varphi^{(\varepsilon_0)}$ and a sequence of independent subordinators $(S^{(\varepsilon_n)})_{n\geq 0}$ such that for each n, $S^{(\varepsilon_n)}$ has Laplace exponent $\Gamma^{(\varepsilon_n)}$. Assume moreover that $X^{(\varepsilon_0)}$ is independent of the sequence $(S^{(\varepsilon_n)})_{n\geq 0}$. Then the sequence of Lévy processes $X^{(\varepsilon_n)} := X^{(\varepsilon_0)} + \sum_{k=0}^{n-1} S^{(\varepsilon_k)}$ satisfies the conditions of the statement.

Note that $\Sigma^{(n)} := \sum_{k=0}^{n-1} S^{(\varepsilon_k)}$ is an increasing sequence of subordinators and that $\Sigma^{(n)}$ has Laplace exponent, $\varphi^{(\varepsilon_n)} - \varphi^{(\varepsilon_0)}$. Moreover, $\lim_{n\to\infty} \varphi^{(\varepsilon_n)} = \varphi$. Therefore, for each $t \geq 0$, $(\Sigma_t^{(n)})$ converges weakly toward an infinitely divisible distribution having $\varphi - \varphi^{(\varepsilon_0)}$ as Laplace exponent. Then for each $t \geq 0$, the a.s. limit $\Sigma_t = \sum_{k=0}^{\infty} S_t^{(\varepsilon_k)}$ of $\Sigma_t^{(n)}$ is a.s. finite and has Laplace exponent $\varphi - \varphi^{(\varepsilon_0)}$. Moreover (Σ_t) defines a càdlàg non decreasing process. It is actually a subordinator with Laplace exponent $\varphi - \varphi^{(\varepsilon_0)}$. Since for all $t \geq 0$, $\sup_{0 \leq s \leq t} |\Sigma_s^{(n)} - \Sigma_s| = |\Sigma_t^{(n)} - \Sigma_t|$, the sequence of subordinators $(\Sigma^{(n)})$ converges uniformly over any closed interval of \mathbb{R}_+ , almost surely, toward the process Σ . This implies the last assertion of Theorem 2.3 regarding the sequence $(X^{(\varepsilon_n)})$.

4.2 Proof of the results in Subsection 3.2 (convergence in distribution)

Recall that $\phi^{(n)} = -\varphi^{(n)}$, $n \ge 0$ and $\phi = -\varphi$ are the branching mechanisms of $Z^{(n)}$, $n \ge 0$ and Z, respectively and that $(u_t^{(n)}(\lambda), t, \lambda \ge 0), n \ge 0$ and $(u_t(\lambda), t, \lambda \ge 0)$ are the corresponding Laplace exponents. Then we first state some of the elementary properties of these Laplace exponents, a part of which can also be found in Chapter 3 of [10].

Lemma 4.1.

- 1. For all $t \ge 0$ and $n \ge 0$, the mappings $\lambda \mapsto u_t^{(n)}(\lambda)$ and $\lambda \mapsto u_t(\lambda)$ are continuous and increasing on $[0, \infty)$.
- 2. For all $\lambda \in (0, \rho)$, the mapping $t \mapsto u_t(\lambda)$ is continuous and increasing on $[0, \infty)$. Moreover it is valued in the set $(0, \rho)$.
- 3. For all $\lambda \in (0, \rho)$, there is $n_{\lambda} \geq 1$ such that for all $n \geq n_{\lambda}$, the mappings $t \mapsto u_t^{(n)}(\lambda)$ are continuous and increasing on $[0, \infty)$. Moreover they are valued in the set $(0, \rho)$.

4. Let ρ_0 be the largest root of $\phi^{(0)}$. Then for all $t \ge 0$ and $\lambda \in (0, \rho_0)$, the sequence $(u_t^{(n)}(\lambda))_{n\ge 0}$ is non decreasing and $\lim_{n\to\infty} u_t^{(n)}(\lambda) = u_t(\lambda)$.

Proof. The first assertion follows directly from the definition (2.1) as well as continuity of the mappings in 2. and 3.

We first prove that for $\lambda \in (0, \rho)$, the mapping $t \mapsto u_t(\lambda)$ is increasing on $[0, \infty)$. From (2.2), $\frac{\partial u_t}{\partial t}(\lambda)|_{t=0} = \phi(u_0(\lambda)) = \phi(\lambda) > 0$, so that $u_{\delta}(\lambda) > u_0(\lambda) = \lambda$, for all $\delta > 0$ small enough. Let t > 0, then from the mean value theorem applied to the function $\lambda \mapsto u_t(\lambda)$, there is $\lambda_{\delta} \in [\lambda, u_{\delta}(\lambda)]$, such that,

$$u_t(u_{\delta}(\lambda)) - u_t(\lambda) = u_{t+\delta}(\lambda) - u_t(\lambda) = \frac{\partial}{\partial \lambda} u_t(\lambda_{\delta})(u_{\delta}(\lambda) - \lambda),$$

where we have used the semigroup property of $u_t(\lambda)$ in the first equality. Note that both $u_{\delta}(\lambda)$ and λ_{δ} tend to λ , as δ tends to 0, hence it follows from the above equality that,

$$\lim_{\delta \to 0} \frac{u_{t+\delta}(\lambda) - u_t(\lambda)}{u_{\delta}(\lambda) - \lambda} = \frac{\partial}{\partial \lambda} u_t(\lambda) > 0.$$

Multiplying and dividing the left hand side of this equality by δ yields,

$$\lim_{\delta \to 0} \frac{u_{t+\delta}(\lambda) - u_t(\lambda)}{u_{\delta}(\lambda) - \lambda} = \frac{\partial}{\partial t} u_t(\lambda) \left(\frac{\partial u_t}{\partial t}(\lambda)|_{t=0}\right)^{-1} > 0$$

Hence, $\frac{\partial}{\partial t}u_t(\lambda)$ has the same sign as $\frac{\partial u_t}{\partial t}(\lambda)|_{t=0} = \phi(\lambda) > 0$. This proves that $t \mapsto u_t(\lambda)$ is increasing on $[0,\infty)$. Then it is clear that of all $\lambda \in (0,\rho)$, $u_t(\lambda) > 0$. Moreover $u_t(\lambda) < \rho$ since $\frac{\partial u_t}{\partial t}(\lambda) = \phi(u_t(\lambda)) > 0$.

Let ρ_n be the largest root of $\phi^{(n)}$, then since $(\phi^{(n)})$ is increasing and $\phi^{(n)} \leq \phi$, the sequence (ρ_n) is increasing and $\rho_n \leq \rho$. Moreover, $\lim_{n\to\infty} \rho_n = \rho$, so that for all $\lambda \in (0, \rho)$, there is $n_{\lambda} \geq 1$ such that for all $n \geq n_{\lambda}$, $\lambda \in (0, \rho_n)$. Then the proof of 3. follows this of 2. along the lines.

Let us emphasize that from 2. and 3., for $n \ge n_{\lambda}$, $\phi^{(n)}$ has no root between $\lambda \in (0, \rho_n)$ and $u_t^{(n)}(\lambda)$ and ϕ has no root between $\lambda \in (0, \rho)$ and $u_t(\lambda)$. Since $\phi^{(n)} \le \phi^{(n+1)}$ for all n, we derive from (2.4) that for all $n \ge 0$, $\lambda \in (0, \rho_n)$ and $t \ge 0$,

$$\int_{\lambda}^{u_t^{(n+1)}(\lambda)} \frac{du}{\phi^{(n+1)}(u)} = t = \int_{\lambda}^{u_t^{(n)}(\lambda)} \frac{du}{\phi^{(n)}(u)} \le \int_{\lambda}^{u_t^{(n+1)}(\lambda)} \frac{du}{\phi^{(n)}(u)}$$

Therefore, $u_t^{(n)}(\lambda) \leq u_t^{(n+1)}(\lambda)$, for all $\lambda \in (0, \rho_n)$. On the other hand, we derive from

$$\int_{\lambda}^{u_t^{(n)}(\lambda)} \frac{du}{\phi(u)} \le \int_{\lambda}^{u_t^{(n)}(\lambda)} \frac{du}{\phi^{(n)}(u)} = t = \int_{\lambda}^{u_t^{(n)}(\lambda)} \frac{du}{\phi(u)} + \int_{u_t^{(n)}(\lambda)}^{u_t(\lambda)} \frac{du}{\phi(u)}$$
(4.1)

that $u_t^{(n)}(\lambda) \leq u_t(\lambda)$, for all $n \geq 0$ and $\lambda \in (0, \rho_n)$. Moreover, since $u_t(\lambda) < \rho$, we derive from (4.1) and dominated convergence theorem that for all $\lambda \in (0, \rho_0)$, $\lim_{n \to \infty} u_t^{(n)}(\lambda) = u_t(\lambda)$.

Recall that $(h(n))_{n\geq 0}$ is a decreasing sequence such that $\lim_{n\to\infty} h(n) = 0$. Moreover, with no loss of generality, we shall henceforth assume that $h(0) < \rho_0$.

Lemma 4.2.

1. For all $t \ge 0$ and $n \ge 0$, $0 < u_t^{(n)}(h(n)) \le u_t(h(0)) < \rho$.

2. Assume that $(u_t^{(n)}(h(n)))_{n\geq 0}$ converges for all $t\geq 0$ and set

$$l_t(h) := \lim_{n \to \infty} u_t^{(n)}(h(n)) \,.$$

Then the mapping $t \mapsto l_t(h)$ is continuous and non decreasing on $[0,\infty)$. In this case, we set

$$c(h) := \sup\{t \ge 0 : l_t(h) = 0\}.$$

3. The sequence $(u_t^{(n)}(h(n)))_{n\geq 0}$ converges for all $t\geq 0$ if and only if $\lim_{n\to\infty} \int_{h(n)}^{\theta} \frac{du}{\phi^{(n)}(u)}$ exists in $[0,\infty]$ for some $\theta \in (0,\rho)$. In this case,

$$\lim_{n \to \infty} \int_{h(n)}^{\theta} \frac{du}{\phi^{(n)}(u)} = c(h) + \int_{0}^{\theta} \frac{du}{\phi(u)}, \quad \text{for all } \theta \in (0, \rho).$$
(4.2)

If $c(h) < \infty$, then for all $t \ge c(h)$,

$$\lim_{n \to \infty} u_t^{(n)}(h(n)) = u_{t-c(h)}(0).$$

Proof. Since (h(n)) is decreasing and $h(0) \in (0, \rho_0)$, from 1. and 4. in Lemma 4.1, for all $t \ge 0, n \ge 0$ and $k \ge n, u_t^{(n)}(h(n)) \le u_t^{(n)}(h(0)) \le u_t^{(k)}(h(0))$. Then by letting k go to ∞ and applying 2. in Lemma 4.1 again, we obtain $u_t^{(n)}(h(n)) \le u_t^{(n)}(h(0)) \le u_t(h(0))$. Since $h(0) \in (0, \rho_0)$ and $\rho_0 < \rho$, assertion 2. in Lemma 4.1 implies that $u_t(h(0)) < \rho$.

Let us now prove assertion 2. Assume that $(u_t^{(n)}(h(n)))_{n\geq 0}$ converges for all $t\geq 0$. Then from 3. in Lemma 4.1, for all $\lambda \in (0, \rho)$ and *n* sufficiently large, the functions $t\mapsto u_t^{(n)}(\lambda)$ are increasing, therefore $t\mapsto l_t(h)$ is non decreasing. Moreover $t\mapsto l_t(h)$ is continuous. Indeed from (2.4) applied to $\phi^{(n)}$, for all $n\geq 0$ and $0\leq s\leq t$,

$$\int_{u_s^{(n)}(h(n))}^{u_t^{(n)}(h(n))} \frac{du}{\phi^{(n)}(u)} = t - s \,,$$

so that from Fatou's lemma and the assumption

$$\int_{l_s(h)}^{l_t(h)} \frac{du}{\phi(u)} \le \liminf_n \int_{u_s^{(n)}(h(n))}^{u_t^{(n)}(h(n))} \frac{du}{\phi^{(n)}(u)} = t - s \,.$$

The continuity of $t \mapsto l_t(h)$ follows from this.

Let us finally prove 3. Assume first that $l_t(h) = \lim_{n \to \infty} u_t^{(n)}(h(n))$ exists for all $t \ge 0$. If $c(h) = \infty$, that is for all $t \ge 0$, $l_t(h) = 0$, then by Fatou's Lemma in

$$\int_{h(n)}^{\theta} \frac{du}{\phi^{(n)}(u)} = t + \int_{u_t^{(n)}(h(n))}^{\theta} \frac{du}{\phi^{(n)}(u)}, \quad \theta \in (0, \rho_n),$$
(4.3)

we obtain, for all $\theta \in (0, \rho)$,

$$\liminf_{n \to \infty} \int_{h(n)}^{\theta} \frac{du}{\phi^{(n)}(u)} \ge t + \int_{0}^{\theta} \frac{du}{\phi(u)}$$

and taking the limit when t goes to infinity implies $\lim_{n\to\infty} \int_{h(n)}^{\theta} \frac{du}{\phi^{(n)}(u)} = \infty$. If $c(h) < \infty$, then let t > c(h), so that $l_t(h) > 0$. Using dominated convergence in (4.3) yields for all

 $\theta \in (0,\rho), \lim_{n \to \infty} \int_{h(n)}^{\theta} \frac{du}{\phi^{(n)}(u)} = t + \int_{l_t(h)}^{\theta} \frac{du}{\phi(u)}$, and letting t tend to c(h) in this equality,

we obtain by continuity of the function $t \mapsto l_t(h)$ that $\lim_{n \to \infty} \int_{h(n)}^{\theta} \frac{du}{\phi^{(n)}(u)} = c(h) + \int_{0}^{\theta} \frac{du}{\phi(u)}$. This implies that,

$$\int_0^{l_t(h)} \frac{du}{\phi(u)} = t - c(h),$$

for all $t \ge c(h)$, that is $l_t(h) = u_{t-c(h)}(0)$.

Conversely, assume that $\lim_{n\to\infty}\int_{h(n)}^{\theta}\frac{du}{\phi^{(n)}(u)}$ exists in $[0,\infty]$ and that for some t>0, there are two subsequences $(v_t^{(n)})$ and $(w_t^{(n)})$ of $(u_t^{(n)}(h(n)))$ which converge toward $v_t > 0$ and 0 respectively. Then from dominated convergence in (4.3) where we replaced $u_t^{(n)}(h(n))$ by $v_t^{(n)}$ we obtain,

$$\lim_{n \to \infty} \int_{h(n)}^{\theta} \frac{du}{\phi^{(n)}(u)} = t + \int_{v_t}^{\theta} \frac{du}{\phi(u)},\tag{4.4}$$

and from Fatou's lemma in (4.3) where we replaced $u_t^{(n)}(h(n))$ by $w_t^{(n)}$ we obtain,

$$\lim_{n \to \infty} \int_{h(n)}^{\theta} \frac{du}{\phi^{(n)}(u)} \ge t + \int_{0}^{\theta} \frac{du}{\phi(u)}.$$

This is contradictory, so either any subsequence of $(u_t^{(n)}(h(n)))$ converges toward a positive limit, and in this case, from (4.4) these limits are identical, or $(u_t^{(n)}(h(n)))$ converges toward 0. If t = 0, then $u_t^{(n)}(h(n)) = h(n) \rightarrow 0 = u_0(0)$. Finally, (4.2) follows by taking t = c(h) in (4.4), where actually $v_t = l_t(h)$ and by recalling that from part 2., $l_{c(h)}(h) = 0.$

Lemma 4.3. For all $t \ge 0$ and $x \ge 0$,

$$\lim_{n \to \infty} Z_t^{(n)} = Z_t \quad and \quad \lim_{n \to \infty} \overline{Z}_t^{(n)} = \overline{Z}_t \,, \quad \mathbb{P}_x \text{-}a.s.$$

Moreover, on the event $\{\zeta \leq t\}$,

$$\lim_{n \to \infty} \frac{Z_t^{(n)}}{\overline{Z}_t^{(n)}} = 1, \quad \mathbb{P}_x \text{-}a.s.$$

Proof. Fix $t \ge 0$, $x \ge 0$ and let (\mathcal{F}_t) be the filtration generated by all the processes $X^{(n)}$, $n \ge 0$. Define also $I_t^{(n)} = \inf \left\{ s : \int_0^s \frac{du}{X_t^{(n)}} > t \right\}$ and $\tau^{(n)} = \inf \{ t : X_t^{(n)} \le 0 \}$. Then for each $t \ge 0$, the sequence $(I_t^{(n)} \land \tau^{(n)})$ is an increasing sequence of stopping times in the filtration (\mathcal{F}_t) , and it satisfies $\lim_n I_t^{(n)} \wedge \tau^{(n)} = I_t \wedge \tau$, \mathbb{P}_x -a.s. Moreover, X is a Lévy process in the filtration (\mathcal{F}_t) and it is quasi-left-continuous. Therefore $\lim_n X(I_t^{(n)} \wedge \tau^{(n)}) =$ $X(I_t \wedge \tau)$, \mathbb{P}_x -a.s. on the set $\{I_t \wedge \tau < \infty\}$, see Proposition 7 of Chapter I in [1]. Then from the construction of $Z^{(n)}$ and Z,

$$|Z_t^{(n)} - Z_t| \leq \sup_{s \leq I_t \wedge \tau} |X_s^{(n)} - X_s| + |X_{I_t^{(n)} \wedge \tau^{(n)}} - X_{I_t \wedge \tau}|, \qquad (4.5)$$

and from Theorem 2.3, $\lim_{n} Z_t^{(n)} = Z_t$, \mathbb{P}_x -a.s. on the set $\{I_t \wedge \tau < \infty\}$. On the set $\{I_t \wedge \tau = \infty\}$, $\lim_{n} I_t^{(n)} \wedge \tau^{(n)} = \infty$, \mathbb{P}_x -a.s., so that $\lim_{n} Z_t^{(n)} = \lim_{n} X^{(n)}(I_t^{(n)} \wedge \tau^{(n)}) =$

 $\infty = Z_t$, \mathbb{P}_x -a.s. The proof of the fact that $\lim_n \overline{Z}_t^{(n)} = \overline{Z}_t = \infty$, \mathbb{P}_x -a.s. follows the same arguments together with the fact that $\sup_{s \leq t} |\overline{X}_s^{(n)} - \overline{X}_s| = \overline{X}_t - \overline{X}_t^{(n)}$, so that $\lim_n \sup_{s \leq t} |\overline{X}_s^{(n)} - \overline{X}_s| = 0$, \mathbb{P} -a.s. for all $t \geq 0$.

Let us now prove the second assertion. First observe that for all $0 \le m \le n$ and $t \ge 0$,

$$\frac{X_t^{(n)}}{\overline{X}_t^{(n)}} \ge \frac{X_t^{(m)}}{\overline{X}_t^{(m)}}.$$
(4.6)

Indeed, write $X_t^{(n)} = X_t^{(m)} + S_t^{m,n}$, where $S^{m,n}$ is a subordinator. Then $\overline{X}_t^{(n)} \leq \overline{X}_t^{(m)} + S_t^{m,n}$, so that

$$\frac{X_t^{(n)}}{\overline{X}_t^{(n)}} - \frac{X_t^{(m)}}{\overline{X}_t^{(m)}} = \frac{\overline{X}_t^{(m)} X_t^{(n)} - \overline{X}_t^{(n)} X_t^{(m)}}{\overline{X}_t^{(n)} \overline{X}_t^{(m)}} = \frac{\overline{X}_t^{(m)} (X_t^{(m)} + S_t^{m,n}) - (\overline{X}_t^{(m)} + S_t^{m,n}) X_t^{(m)}}{\overline{X}_t^{(n)} \overline{X}_t^{(m)}} \\ \ge \frac{S_t^{m,n} (\overline{X}_t^{(m)} - X_t^{(m)})}{\overline{X}_t^{(n)} \overline{X}_t^{(m)}} \ge 0.$$

Recall from Subsection 3.1 that we choose ε_0 small enough so that $0 < \phi^{(n)'}(0) = \phi'(\varepsilon_n) < \infty$, for all $n \ge 0$, and hence $\mathbb{E}(X_1^{(n)}) > 0$, for all $n \ge 0$. Let $\delta \in (0, 1)$, then from Lemma 2.1 in [9], there exists $t_0 > 0$ such that \mathbb{P} -a.s., for all $t \ge t_0$, $X_t^{(0)}/\overline{X}_t^{(0)} \ge 1 - \delta$ and then from (4.6) for all $n \ge 0$, $X_t^{(n)}/\overline{X}_t^{(n)} \ge 1 - \delta$. On the other hand, note that for all $t \ge 0$, $\{\zeta \le t\} \subset \{I_t \land \tau = \infty\}$ and hence $\lim_n I_t^{(n)} \land \tau^{(n)} = \infty$, \mathbb{P} -a.s. on the set $\{\zeta \le t\}$. This yields,

$$\lim_{n} \frac{Z_{t}^{(n)}}{\overline{Z}_{t}^{(n)}} = \lim_{n} \frac{X^{(n)}(I_{t}^{(n)} \wedge \tau^{(n)})}{\overline{X}^{(n)}(I_{t}^{(n)} \wedge \tau^{(n)})} \ge 1 - \delta, \quad \mathbb{P}\text{-a.s. on the set } \{\zeta \le t\},$$

which allows us to conclude, δ being arbitrary.

Lemma 4.4. Let e be some exponentially distributed random variable with parameter 1 and assume that e is independent of the family $(Z^{(n)}, n \ge 0)$. Fix t > 0 and x > 0.

- 1. If $\lim_{n \to \infty} u_t^{(n)}(h(n))$ exists, then $\lim_{n \to \infty} u_t^{(n)}(kh(n)) = \lim_{n \to \infty} u_t^{(n)}(h(n))$, for all k > 0.
- 2. If $\lim_{n\to\infty} \mathbb{P}_x(\overline{Z}_t^{(n)} \le e/h(n)) = e^{-xu_t(0)}$, then $\lim_{n\to\infty} \mathbb{P}_x(Z_t^{(n)} \le e/h(n)) = e^{-xu_t(0)}$.
- 3. $\lim_{n\to\infty} u_t^{(n)}(h(n))$ exists if and only if $\lim_{n\to\infty} \mathbb{P}_x(\overline{Z}_t^{(n)} \leq ke/h(n))$ exists for all k > 0 and does not depend on k. In this case, for all k > 0, $\lim_{n\to\infty} \mathbb{P}_x(\overline{Z}_t^{(n)} \leq ke/h(n)) = e^{-xl_t(h)}$, where $l_t(h) := \lim_{n\to\infty} u_t^{(n)}(h(n))$.

Proof. Let $\theta \in (0, \rho_0)$, so that none of the functions $\phi^{(n)}$ vanishes on $(0, \theta)$. Recall also that $h(0) < \rho_0$. Let k > 0 and n such that $kh(n) < \theta$ and write

$$\int_{h(n)}^{kh(n)} \frac{du}{\phi^{(n)}(u)} = \int_{h(n)}^{\theta} \frac{du}{\phi^{(n)}(u)} - \int_{kh(n)}^{\theta} \frac{du}{\phi^{(n)}(u)}$$

Since $\phi^{(n)}$ is increasing and differentiable,

$$\left| \int_{h(n)}^{kh(n)} \frac{du}{\phi^{(n)}(u)} \right| \le \frac{|k-1|h(n)|}{\phi^{(n)}((k\wedge 1)h(n))} = \frac{|k-1|}{(k\wedge 1)\phi'(\alpha_n)},$$

where $\alpha_n \in (\varepsilon_n, \varepsilon_n + (k \wedge 1)h(n))$. Since $\phi'(0) = \infty$, it follows from the above inequality that $\lim_{n\to\infty} \int_{h(n)}^{kh(n)} \frac{du}{\phi^{(n)}(u)} = 0$, so that if one of the sequences $(\int_{h(n)}^{\theta} \frac{du}{\phi^{(n)}(u)})$ or $(\int_{kh(n)}^{\theta} \frac{du}{\phi^{(n)}(u)})$ converges in $[0, \infty]$ as *n* tends to infinity, then both converge and they have the same limit. Then part 1. follows from part 3. of Lemma 4.2.

Before proving 2. and 3., recall from Lemma 4.3 that $Z_t^{(n)}$ and $\overline{Z}_t^{(n)}$ converge \mathbb{P}_x -almost surely to Z_t and \overline{Z}_t , respectively. Therefore, for all k > 0,

$$\lim_{n \to \infty} \mathbb{P}_x(Z_t^{(n)} \le \operatorname{ke}/h(n), t < \zeta) = \mathbb{P}_x(Z_t < \infty, t < \zeta) = \mathbb{P}_x(t < \zeta) = e^{-xu_t(0)},$$
$$\lim_{n \to \infty} \mathbb{P}_x(\overline{Z}_t^{(n)} \le \operatorname{ke}/h(n), t < \zeta) = \mathbb{P}_x(\overline{Z}_t < \infty, t < \zeta) = \mathbb{P}_x(t < \zeta) = e^{-xu_t(0)}.$$

Then let us prove assertion 2. and assume that $\lim_{n\to\infty} \mathbb{P}_x(\overline{Z}_t^{(n)} \leq e/h(n)) = e^{-xu_t(0)}$. From (4.7) this means that $\lim_{n\to\infty} \mathbb{P}_x(\overline{Z}_t^{(n)} \leq e/h(n), t \geq \zeta) = 0$. Then let a < 1 and write

$$\mathbb{P}_x\left(a^{-1}Z_t^{(n)} \le e/h(n), t \ge \zeta\right) \le \mathbb{P}_x\left(\overline{Z}_t^{(n)} \le e/h(n), Z_t^{(n)}/\overline{Z}_t^{(n)} \in [a, 1], t \ge \zeta\right) \\ +\mathbb{P}_x\left(Z_t^{(n)}/\overline{Z}_t^{(n)} \in [a, 1]^c, t \ge \zeta\right).$$

This inequality together with Lemma 4.3 implies that $\lim_{n\to\infty} \mathbb{P}_x \left(a^{-1} Z_t^{(n)} \leq e/h(n), t \geq \zeta \right) = 0$, so that from (4.7), $\lim_{n\to\infty} \mathbb{P}_x \left(a^{-1} Z_t^{(n)} \leq e/h(n) \right) = e^{-xu_t(0)}$. But $\mathbb{P}_x \left(a^{-1} Z_t^{(n)} \leq e/h(n) \right) = \mathbb{E}_x (e^{-a^{-1}h(n)Z_t^{(n)}}) = e^{-xu_t^{(n)}(a^{-1}h(n))}$, and the conclusion follows from part 1. of this lemma.

 $\mathbb{E}_x(e^{-a^{-1}h(n)Z_t^{(n)}}) = e^{-xu_t^{(n)}(a^{-1}h(n))}, \text{ and the conclusion follows from part 1. of this lemma.}$ Finally let us prove 3. Assume that $\lim_{n\to\infty} \mathbb{P}_x(\overline{Z}_t^{(n)} \leq ke/h(n))$ exists for all k > 0and does not depend on k. Then from (4.7) $\lim_{n\to\infty} \mathbb{P}_x(\overline{Z}_t^{(n)} \leq ke/h(n), t \geq \zeta)$ exists for all k > 0 and is equal to $\lim_{n\to\infty} \mathbb{P}_x(\overline{Z}_t^{(n)} \leq e/h(n), t \geq \zeta)$. Now let a > 1 and write

$$\mathbb{P}_x\left(aZ_t^{(n)} \le e/h(n), \,\overline{Z}_t^{(n)}/Z_t^{(n)} \in [1,a], t \ge \zeta\right) \le \mathbb{P}_x\left(\overline{Z}_t^{(n)} \le e/h(n), \, t \ge \zeta\right) \,, \tag{4.8}$$

so that from Lemma 4.3,

$$\limsup_{n \to \infty} \mathbb{P}_x \left(a Z_t^{(n)} \le e/h(n), \ t \ge \zeta \right) \le \lim_{n \to \infty} \mathbb{P}_x \left(\overline{Z}_t^{(n)} \le e/h(n), \ t \ge \zeta \right) .$$
(4.9)

But since $\lim_{n\to\infty} \mathbb{P}_x(\overline{Z}_t^{(n)} \le ke/h(n), t \ge \zeta)$ does not depend on k, replacing h by h/a in (4.9), we can write

$$\limsup_{n \to \infty} \mathbb{P}_x \left(Z_t^{(n)} \le \mathbf{e}/h(n), \ t \ge \zeta \right) \le \lim_{n \to \infty} \mathbb{P}_x \left(\overline{Z}_t^{(n)} \le \mathbf{e}/h(n), \ t \ge \zeta \right) \,.$$

Moreover, the inequality

$$\lim_{n \to \infty} \mathbb{P}_x \left(\overline{Z}_t^{(n)} \le \mathbf{e}/h(n), \ t \ge \zeta \right) \le \liminf_{n \to \infty} \mathbb{P}_x \left(Z_t^{(n)} \le \mathbf{e}/h(n), \ t \ge \zeta \right)$$

is straightforward. So we have proved that $\lim_{n\to\infty} \mathbb{P}_x\left(Z_t^{(n)} \leq e/h(n), t \geq \zeta\right)$ exists. We conclude from the equality,

$$\mathbb{P}_x\left(Z_t^{(n)} \le e/h(n)\right) = e^{-xu_t^{(n)}(h(n))}.$$
(4.10)

Conversely assume that $\lim_{n\to\infty} u_t^{(n)}(h(n))$ exists. Then from part 1. of the present lemma, $\lim_{n\to\infty} u_t^{(n)}(kh(n)) = \lim_{n\to\infty} u_t^{(n)}(h(n))$, exists for all k > 0 and therefore

 $\lim_{n} \mathbb{P}_{x} \left(Z_{t}^{(n)} \leq e/kh(n) \right) = \lim_{n} e^{-xu_{t}^{(n)}(kh(n))} \text{ exists for all } k > 0. \text{ Then we derive from}$ (4.7) that $\lim_{n \to \infty} \mathbb{P}_{x}(Z_{t}^{(n)} \leq ke/h(n), t \geq \zeta)$ exists for all k > 0 and does not depend on k. So from the same argument using Lemma 4.3, developed in (4.8) and (4.9), for all k > 0,

$$\lim_{n \to \infty} \mathbb{P}_x \left(Z_t^{(n)} \le e/h(n), t \ge \zeta \right) \le \liminf_{n \to \infty} \mathbb{P}_x \left(\overline{Z}_t^{(n)} \le ke/h(n), t \ge \zeta \right) \,.$$

We conclude from the inequalities,

$$\limsup_{n \to \infty} \mathbb{P}_x \left(\overline{Z}_t^{(n)} \le k e/h(n), t \ge \zeta \right) \le \lim_{n \to \infty} \mathbb{P}_x \left(Z_t^{(n)} \le k e/h(n), t \ge \zeta \right)$$
$$= \lim_{n \to \infty} \mathbb{P}_x \left(Z_t^{(n)} \le e/h(n), t \ge \zeta \right)$$

and the equality (4.10).

Lemma 4.5. Fix x, t > 0, then $\lim_{n\to\infty} u_t^{(n)}(h(n))$ exists if and only if $\lim_{n\to\infty} \mathbb{P}_x(\sigma_{k/h(n)}^{(n)} > t)$ exists for all k > 0 and does not depend on k. In this case, $\lim_{n\to\infty} \mathbb{P}_x(\sigma_{k/h(n)}^{(n)} > t) = e^{-xl_t(h)}$, where $\lim_{n\to\infty} u_t^{(n)}(h(n)) := l_t(h)$.

Proof. Let e be as in Lemma 4.4 and note that $z \mapsto \mathbb{P}_x(\sigma_{z/h(n)}^{(n)} > t)$ is increasing. Then on the one hand, for all $\delta > 0$,

$$\mathbb{P}_{x}(\sigma_{\delta/h(n)}^{(n)} > t)e^{-\delta} \leq \int_{\delta}^{\infty} \mathbb{P}_{x}(\sigma_{z/h(n)}^{(n)} > t)e^{-z} dz \\
\leq \int_{0}^{\infty} \mathbb{P}_{x}(\sigma_{z/h(n)}^{(n)} > t)e^{-z} dz = \mathbb{P}_{x}(\sigma_{e/h(n)}^{(n)} > t), \quad (4.11)$$

and on the other hand, for all d > 0,

$$\mathbb{P}_{x}(\sigma_{e/h(n)}^{(n)} > t) \leq \mathbb{P}_{x}\left(\sigma_{e/h(n)}^{(n)} > t, e/h(n) < d/h(n)\right) + \mathbb{P}_{x}\left(e/h(n) > d/h(n)\right) \\
= \int_{0}^{d} \mathbb{P}_{x}(\sigma_{z/h(n)}^{(n)} > t)e^{-z} dz + \int_{d}^{\infty} e^{-z} dz \\
\leq (1 - e^{-d})\mathbb{P}_{x}(\sigma_{d/h(n)}^{(n)} > t) + e^{-d}.$$
(4.12)

Then assume that $\lim_{n\to\infty} u_t^{(n)}(h(n))$ exists and set $\lim_{n\to\infty} u_t^{(n)}(h(n)) := l_t(h)$. Recall that e is as in Lemma 4.4 and note that $\mathbb{P}_x(\sigma_{e/h(n)}^{(n)} > t) = \mathbb{P}_x(\overline{Z}_t^{(n)} < e/h(n))$. Then by part 3. of Lemma 4.4, for all k > 0,

$$\lim_{n \to \infty} \mathbb{P}_x(\sigma_{\mathbf{e}/h(n)}^{(n)} > t) = \lim_{n \to \infty} \mathbb{P}_x(\sigma_{k\mathbf{e}/h(n)}^{(n)} > t) = e^{-xl_t(h)} .$$

$$(4.13)$$

Thanks to (4.13), replacing h by kh/δ (resp. by kh/d) in (4.11) (resp. in (4.12)), we obtain that for all $\delta, d, k > 0$,

$$e^{-\delta} \limsup_{n \to 0} \mathbb{P}_x(\sigma_{k/h(n)}^{(n)} > t) \le e^{-xl_t(h)} \le (1 - e^{-d}) \liminf_{n \to 0} \mathbb{P}_x(\sigma_{k/h(n)}^{(n)} > t) + e^{-d}$$

Taking δ to 0 and d to ∞ shows that $\lim_{n\to\infty} \mathbb{P}_x(\sigma_{k/h(n)}^{(n)} > t) = e^{-xl_t(h)}$.

Conversely if $\lim_{n\to\infty} \mathbb{P}_x(\sigma_{k/h(n)}^{(n)} > t)$ exists for all k > 0 and does not depend on k, then replacing h by h/k in (4.11) and (4.12), we obtain,

$$\lim_{n \to \infty} \mathbb{P}_x(\sigma_{1/h(n)}^{(n)} > t)e^{-\delta} \leq \liminf_{n \to \infty} \mathbb{P}_x(\sigma_{ke/h(n)}^{(n)} > t) \leq \limsup_{n \to \infty} \mathbb{P}_x(\sigma_{ke/h(n)}^{(n)} > t)$$
$$\leq (1 - e^{-d}) \lim_{n \to \infty} \mathbb{P}_x(\sigma_{1/h(n)}^{(n)} > t) + e^{-d}.$$

Taking δ to 0 and d to ∞ shows that $\lim_{n\to\infty} \mathbb{P}_x(\sigma_{ke/h(n)}^{(n)} > t)$ exists for all k > 0 and does not depend on k. We conclude from part 2. of Lemma 4.4.

Proof of Theorem 3.1. Equivalence between 1. and 3. is given by Lemma 4.5. Equivalence between 2. and 3. is given by 3. in Lemma 4.2. Then still from 3. in Lemma 4.2 and Lemma 4.5, for all $x, t \ge 0$, $\lim_{n\to\infty} \mathbb{P}_x(\sigma_{1/h(n)}^{(n)} > t) = \lim_{n\to\infty} e^{-xu_t^{(n)}(h(n))} = e^{-u_{(t-c(h))}+(0)} = \mathbb{P}_x(\zeta + c(h) > t)$, which achieves the proof of the theorem.

Proof of Proposition 3.1. This proof follows from 2. and 3. in Lemma 4.4 and the fact that $\mathbb{P}_x(Z_t^{(n)} \leq e/h(n)) = e^{-xu_t^{(n)}(h(n))}$.

Proof of Proposition 3.2. Assume that $\sigma_{1/h(n)}^{(n)} \xrightarrow{(d)}{n \to \infty} \zeta$, that is, for all t > 0, $\lim_{n \to \infty} \mathbb{P}_x(\overline{Z}_t^{(n)} \le 1/h(n)) = e^{-xu_t(0)}$. Note that we can replace the exponential time e by 1 in (4.7) so that for all decreasing sequence (f(n)) such that $\lim_{n \to \infty} f(n) = 0$,

$$\lim_{n \to \infty} \mathbb{P}_x(\overline{Z}_t^{(n)} \le 1/f(n), t < \zeta) = \mathbb{P}_x(\overline{Z}_t < \infty, t < \zeta) = \mathbb{P}_x(t < \zeta) = e^{-xu_t(0)}.$$
 (4.14)

From the assumption and (4.14) applied to f = h, $\lim_{n\to\infty} \mathbb{P}_x(\overline{Z}_t^{(n)} \leq 1/h(n), t \geq \zeta) = 0$. Moreover, since $h(n) \leq \tilde{h}(n)$, for all *n* large enough,

$$\mathbb{P}_x(\overline{Z}_t^{(n)} \le 1/\tilde{h}(n), t \ge \zeta) \le \mathbb{P}_x(\overline{Z}_t^{(n)} \le 1/h(n), t \ge \zeta)$$

so that the left hand side of this inequality tends to 0 when n goes to infinity. Applying this together with (4.14) for $f = \tilde{h}$ gives $\sigma_{1/\tilde{h}(n)}^{(n)} \xrightarrow{(d)}{n \to \infty} \zeta$.

Proof of Proposition 3.3. Let (h(n)) be such a sequence. Then there are $k_1, k_2 > 0$ such that for all n sufficiently large, $k_1h(n) \leq \tilde{h}(n) \leq k_2h(n)$, so that from 4. in Lemma 4.1, for all $t \geq 0$, $u_t^{(n)}(k_1h(n)) \leq u_t^{(n)}(\tilde{h}(n)) \leq u_t^{(n)}(k_2h(n))$. Then assertion 1. follows from Theorem 3.1. To prove the second assertion it suffices to note that $\lim_{n\to\infty} \int_{h(n)}^{\tilde{h}(n)} \frac{du}{\phi^{(n)}(u)} = c(h) - c(\tilde{h}) \geq 0$. Then to prove part 3. let us write, for $\theta \in (0, \rho)$,

$$\int_{h(n)}^{\theta} \frac{du}{\phi^{(n)}(u)} - \int_{h(n)}^{\tilde{h}(n)} \frac{du}{\phi^{(n)}(u)} = \int_{\tilde{h}(n)}^{\theta} \frac{du}{\phi^{(n)}(u)}, \qquad (4.15)$$

so that taking the liminf on each side gives,

$$\lim_{n \to \infty} \int_{h(n)}^{\theta} \frac{du}{\phi^{(n)}(u)} - \limsup_{n \to \infty} \int_{h(n)}^{h(n)} \frac{du}{\phi^{(n)}(u)} = \liminf_{n \to \infty} \int_{\tilde{h}(n)}^{\theta} \frac{du}{\phi^{(n)}(u)}$$

But $h \in \mathbb{Z}_0$, so that $\lim_{n\to\infty} \int_{h(n)}^{\theta} \frac{du}{\phi^{(n)}(u)} = \int_0^{\theta} \frac{du}{\phi^{(u)}}$, moreover from Fatou's lemma, $\lim_{n\to\infty} \int_{\tilde{h}(n)}^{\theta} \frac{du}{\phi^{(n)}(u)} \ge \int_0^{\theta} \frac{du}{\phi^{(u)}}$, which shows that $\lim_{n\to\infty} \int_{h(n)}^{\tilde{h}(n)} \frac{du}{\phi^{(n)}(u)} = 0$ and finally from (4.15), $\lim_{n\to\infty} \int_{\tilde{h}(n)}^{\theta} \frac{du}{\phi^{(n)}(u)} = \int_0^{\theta} \frac{du}{\phi^{(u)}}$. We conclude from Theorem 3.1. For the last part, note that since $c(h) = \infty$, for all $t \ge 0$, $l_t(h) = 0$. Then recall from Lemma 4.1 that $\lambda \mapsto u_t^{(n)}(\lambda)$ is increasing so that for all $n, u_t^{(n)}(\tilde{h}(n)) \le u_t^{(n)}(h(n))$, which implies that for all $t \ge 0, l_t(\tilde{h}) = 0$ so that $\tilde{h} \in \mathbb{Z}_{\infty}$.

Let us note the following inequalities which are direct consequences of the concavity of ϕ . Let γ be the unique value such that $\gamma \in (0, \rho)$ and $\phi'(\gamma) = 0$ if X is not a subordinator

and $\gamma = \infty$ otherwise. Then for all $u \in (0, \gamma)$ and $n \ge 0$ such that $u + \varepsilon_n \in (0, \gamma)$, $\phi'(u + \varepsilon_n) \le \frac{\phi^{(n)}(u)}{u} \le \phi'(\varepsilon_n)$, so that

$$\frac{1}{u\phi'(\varepsilon_n)} \le \frac{1}{\phi^{(n)}(u)} \le \frac{1}{u\phi'(u+\varepsilon_n)}.$$
(4.16)

Lemma 4.6.

1. If
$$\lim_{n \to \infty} \int_{h(n)}^{\theta} \frac{du}{\phi^{(n)}(u)} = \int_{0}^{\theta} \frac{du}{\phi(u)}$$
, for some $\theta \in (0, \rho)$, then $\lim_{n \to \infty} \frac{|\log h(n)|}{\phi'(\varepsilon_n)} = 0$.

2. If
$$\lim_{n \to \infty} \frac{|\log h(n)|}{\phi'(\varepsilon_n)} = \infty$$
, then $\lim_{n \to \infty} \int_{h(n)}^{\varepsilon} \frac{du}{\phi^{(n)}(u)} = \infty$ for all $\theta \in (0, \rho)$.

Proof. From the assumption and dominated convergence on $[a, \theta)$, we obtain that for all $a \in (0, \theta)$,

$$\lim_{n \to \infty} \int_{h(n)}^{a} \frac{du}{\phi^{(n)}(u)} = \int_{0}^{a} \frac{du}{\phi(u)}$$

On the other hand recall that $\lim_{n\to\infty} \phi'(\varepsilon_n) = \infty$. Then for all $a \in (0, \gamma)$ and $h(n) \leq a$, (4.16) yields,

$$\int_{h(n)}^{a} \frac{du}{\phi^{(n)}(u)} \ge \frac{\log(a)}{\phi'(\varepsilon_n)} - \frac{\log(h(n))}{\phi'(\varepsilon_n)} \implies \limsup_{n \to \infty} \frac{|\log(h(n))|}{\phi'(\varepsilon_n)} \le \int_{0}^{a} \frac{du}{\phi(u)}$$

The first assertion is obtained by taking the limit when a tends to 0 in the last inequality. Then second assertion is a direct consequence of (4.16).

Proof of Proposition 3.4. It follows directly from Lemma 4.6 and the definitions of \mathcal{Z} and \mathcal{Z}_0 .

Proof of Theorem 3.2. This proof requires the following additional lemma.

Lemma 4.7. If $\liminf_{n\to\infty} \phi(h(n) + \varepsilon_n)/\phi(\varepsilon_n) > 1$, then for all $\theta \in (0, \rho)$,

$$\lim_{n \to \infty} \int_{h(n)}^{\theta} \frac{du}{\phi^{(n)}(u)} = \int_0^{\theta} \frac{du}{\phi(u)},$$

that is $h \in \mathcal{Z}_0$.

Proof. We can assume with no loss of generality that $\theta \in (0, \gamma)$. Then we derive from the monotone convergence theorem that $\lim_{n \to \infty} \int_{h(n)}^{\theta} \frac{du}{\phi(u + \varepsilon_n)} = \int_{0}^{\theta} \frac{du}{\phi(u)}$. On the other hand, for all $n \ge 0$ such that $u + \varepsilon_n \in (0, \theta)$,

$$0 \leq \frac{1}{\phi^{(n)}(u)} - \frac{1}{\phi(u+\varepsilon_n)} = \frac{\phi(\varepsilon_n)}{\phi^{(n)}(u)\phi(u+\varepsilon_n)} \leq \frac{\phi(\varepsilon_n)}{\phi^{(n)}(u)\phi(u)}.$$

Moreover, for $u \ge h(n)$,

$$\frac{\phi(\varepsilon_n)}{\phi^{(n)}(u)} \le \frac{\phi(\varepsilon_n)}{\phi(h(n) + \varepsilon_n) - \phi(\varepsilon_n)} = \frac{1}{\phi(h(n) + \varepsilon_n)/\phi(\varepsilon_n) - 1}$$
(4.17)

and from our assumption, there exists C > 0 such that for all n large enough

$$\mathbb{1}_{[h(n),\theta]}(u)\left(\frac{1}{\phi^{(n)}(u)} - \frac{1}{\phi(u+\varepsilon_n)}\right) \leq \frac{C}{\phi(u)}.$$

Then the result follows from dominated convergence.

We are now ready to achieve the proof of Theorem 3.2. Assume now that $\phi = -\varphi$ is regularly varying at 0, with index $\alpha \in (0,1)$. Then from Theorem 1.7.2b in [2], ϕ' is regularly varying at 0, with index $\alpha - 1$.

We assume first that for all k > 0, $\sigma_{k/h(n)}^{(n)} \xrightarrow{(d)} \zeta + c(h)$, that is $h \in \mathbb{Z}$. If c(h) = 0then by Proposition 3.4, $\lim_{n\to\infty} \frac{|\log h(n)|}{\phi'(\varepsilon_n)} = 0$. Now assume that c(h) > 0, let d > 0 and write,

$$\int_{h(n)}^{\theta} \frac{du}{\phi^{(n)}(u)} = \int_{h(n)}^{d\varepsilon_n} \frac{du}{\phi^{(n)}(u)} + \int_{d\varepsilon_n}^{\theta} \frac{du}{\phi^{(n)}(u)}.$$
(4.18)

Since $\lim_{n} \phi((1+d)\varepsilon_n)/\phi(\varepsilon_n) = (1+d)^{\alpha} > 1$, Lemma 4.7 implies that the second term of the right hand side of (4.18) tends to $\int_0^\theta \frac{du}{\phi(u)}$ as $n \to \infty$, that is, $n \mapsto d\varepsilon_n \in \mathbb{Z}_0$. It follows from Theorem 3.1 that

$$\lim_{n \to \infty} \int_{h(n)}^{d\varepsilon_n} \frac{du}{\phi^{(n)}(u)} = \lim_{n \to \infty} \left(\int_{h(n)}^{\theta} \frac{du}{\phi^{(n)}(u)} - \int_{d\varepsilon_n}^{\theta} \frac{du}{\phi^{(n)}(u)} \right) = c(h) \,.$$

Since c(h) > 0, for all n large enough, $h(n) < d\varepsilon_n$, and then the inequalities (4.16) yield,

$$\frac{\log d\varepsilon_n - \log h(n)}{\phi'(\varepsilon_n)} \le \int_{h(n)}^{d\varepsilon_n} \frac{du}{\phi^{(n)}(u)} \le \frac{\log d\varepsilon_n - \log h(n)}{\phi'(d\varepsilon_n + \varepsilon_n)}.$$
(4.19)

Moreover, since ϕ' is regularly varying,

$$\limsup_{n \to \infty} \frac{-\log h(n)}{\phi'(\varepsilon_n)} \le c(h) \le (d+1)^{1-\alpha} \liminf_{n \to \infty} \frac{-\log h(n)}{\phi'(\varepsilon_n)}.$$

We conclude that 2. implies 1. by taking the limit in this inequality as d tends to 0.

Now suppose that $c(h) = \lim_{n \to \infty} \frac{|\log h(n)|}{\phi'(\varepsilon_n)}$ exists. Assume first that c(h) > 0 and recall that since ϕ' is regularly varying with index $\alpha - 1$, $\lim_{n \to \infty} \frac{|\log \varepsilon_n|}{\phi'(\varepsilon_n)} = 0$. This implies that for all d > 0 and for all n large enough, $h(n) < d\varepsilon_n$. Using (4.18) and (4.19) and taking the limit when d goes to 0 yields $\lim_{n \to \infty} \int_{h(n)}^{\theta} \frac{du}{\phi^{(n)}(u)} = \int_{0}^{\theta} \frac{du}{\phi(u)} + c(h)$. Then Theorem 3.1 allows us to derive that for all k > 0, $\sigma_{k/h(n)}^{(n)} \frac{(d)}{n \to \infty} \zeta + c(h)$. If c(h) = 0, then define the two subsequences (n_i) and (m_i) such that for all $i \ge 0$, $h(n_i) < \varepsilon_{n_i}$ and $h(m_i) \ge \varepsilon_{m_i}$. For the subsequence (n_i) , inequalities (4.18) and (4.19) with d = 1 allow us to obtain that $\lim_{n \to \infty} \int_{h(n)}^{\theta} \frac{du}{\phi^{(n)}(u)} = \int_{0}^{\theta} \frac{du}{\phi(u)}$. We conclude from Theorem 3.1 that 2. holds for the subsequence (n_i) . For the subsequence $(m_i)_{i\in\mathbb{N}}$, since, as already noticed above, $m_i \mapsto \varepsilon_{m_i} \in \mathbb{Z}_0$, it suffices to apply part 3. of Proposition 3.3. Then we have proved that 1. and 2. are equivalent in Theorem 3.2.

It is enough to prove that 3 implies 1. Assume without loss of generality that 3. holds for k = 1. Since for all k > 0 and all n sufficiently large, $kh(n) \le \sqrt{h(n)}$, Proposition 3.2 implies that $\sigma_{k/\sqrt{h(n)}}^{(n)} \xrightarrow{(d)} \zeta$, for all k > 0. Therefore, from the equivalence between parts 1. and 2. of Theorem 3.2, $\lim_{n \to \infty} \frac{|\log \sqrt{h(n)}|}{\phi'(\varepsilon_n)} = 0$ and hence $\lim_{n \to \infty} \frac{|\log h(n)|}{\phi'(\varepsilon_n)} = 0$, which is part 1. of Theorem 3.2.

Proof of the results in Subsection 3.3 4.3

4.3.1Convergence in L_1

Proof of Proposition 3.5. Denote by $\mathbb{P}^{\uparrow}_{x}$ the probability under which X starts from x and is conditioned to never reach 0. Recall that $\mathbb{P}_x(Z_t < \infty) = \mathbb{P}_x(\zeta > t) = e^{-xu_t(0)}$ and the notation $\tau = \inf\{t \ge 0 : X_t \le 0\}$. Note also that for all $t \ge 0$, $\{\tau < \infty\} \subset \{Z_t < \infty\}$ and that for all x > 0, $\mathbb{P}_x(\tau < \infty) = e^{-\rho x}$. Then we can write

$$\mathbb{P}_x^{\uparrow}(\zeta > t) = \mathbb{P}_x(Z_t < \infty, \tau = \infty) / \mathbb{P}_x(\tau = \infty)$$
$$= \left(\mathbb{P}_x(Z_t < \infty) - \mathbb{P}_x(Z_t < \infty, \tau < \infty)\right) / \mathbb{P}_x(\tau = \infty) = \frac{e^{-xu_t(0)} - e^{-\rho x}}{1 - e^{-\rho x}}.$$

Therefore, from (2.2), under \mathbb{P}_x^{\uparrow} , the law of ζ is absolutely continuous on $[0,\infty)$ with respect to the Lebesgue measure and its density is given by $f_x(t) := \frac{x\phi(u_t(0))e^{-xu_t(0)'}}{1-e^{-\rho x}}$. This allows us to write

$$\mathbb{E}_x^{\uparrow}(\zeta^n \mathbb{1}_{\{\zeta < \infty\}}) = \int_0^\infty t^n \frac{x\phi(u_t(0))e^{-xu_t(0)}}{1 - e^{-\rho x}} dt$$

The result follows using the substitution $u_t(0) = \lambda$ and the fact that $u_t(0) = F^{-1}(t)$, where $F: y \mapsto \int_0^y du/\phi(u)$ is a bijection from $[0,\rho)$ to $[0,\infty)$, together with the equality $\mathbb{E}_x(\zeta^n \mathbb{1}_{\{\zeta < \infty\}}) = (1 - e^{-\rho x}) \mathbb{E}_x^{\uparrow}(\zeta^n \mathbb{1}_{\{\zeta < \infty\}}). \text{ Since } |\phi'(\rho)| \in (0, \infty), \ \phi(u) \sim (\rho - u) |\phi'(\rho)| \text{ as}$ $u \uparrow \rho$, so that $F(y) =_{y \to \rho} O(\log(\rho - y))$, which implies that the moments of ζ are finite.

We show the second equality from same arguments.

Recall from Subsection 3.1 the definition of (ε_n) , (h(n)), $(X^{(n)})$ and X. Let us also emphasize that

$$h \in \mathcal{Z}_0$$

throughout the rest of this subsection. Moreover, in this subsection, we assume that X is not a subordinator, in particular $\rho \in (0, \infty)$. Indeed, the proof of Theorem 3.3 is actually simpler in the case where X is a subordinator and we allow ourself to skip it. We also recall the notation $\tau = \inf\{t : X_t \leq 0\}$ and introduce the following one,

$$\tau^{(n)} = \inf\{t : X_t^{(n)} \le 0\}.$$

Lemma 4.8. Let $\psi: [0, \infty) \to [\rho, \infty)$ be the inverse of φ , that is, $\varphi \circ \psi(\lambda) = \lambda$ and define $\psi^{(n)}: [0,\infty) \to [\rho_n,\infty)$ the inverse of $\varphi^{(n)}$ in the same way, where ρ_n is the largest root of $\varphi^{(n)}$. Let $e_n := e/h(n)$ where e is an exponentially distributed random variable with parameter 1 which is independent of $(X^{(n)})$ and X. Then

$$\mathbb{E}_x\left(\int_0^{\mathbf{e}_n} \frac{du}{X_u} \mathbb{I}_{\{\tau=\infty\}}\right) = \int_0^\rho \frac{e^{-\lambda x} - e^{\psi(h(n))x}}{h(n) + \phi(\lambda)} d\lambda, \qquad (4.20)$$

$$\mathbb{E}_{x}\left(\int_{0}^{e_{n}}\frac{du}{X_{u}^{(n)}}\mathbb{1}_{\{\tau^{(n)}=\infty\}}\right) = \int_{0}^{\rho_{n}}\frac{e^{-\lambda x} - e^{-\psi^{(n)}(h(n))x}}{h(n) + \phi^{(n)}(\lambda)}d\lambda.$$
(4.21)

Proof. Let us set $\underline{X}_u = \inf_{t \leq u} X_t$, then

$$\begin{split} \mathbb{E}_{x}(e^{-\lambda X_{u}}\mathbb{1}_{\{\tau=\infty\}}) &= e^{-\lambda x}E(e^{-\lambda X_{u}}\mathbb{1}_{\{\underline{X}_{u}\geq -x\}}P_{X_{u}}(\tau_{-x}=\infty)) \\ &= e^{-\lambda x}E(e^{-\lambda X_{u}}\mathbb{1}_{\{\underline{X}_{u}\geq -x\}}P_{X_{u}}(-\underline{X}_{\infty}\leq x)) \\ &= e^{-\lambda x}E(e^{-\lambda X_{u}}\mathbb{1}_{\{\underline{X}_{u}\geq -x\}}(1-e^{-\rho(X_{u}+x)}))\,, \end{split}$$

since $-\underline{X}_{\infty}$ is exponentially distributed with parameter ρ . Let $\varepsilon > 0$ and set $\mathbf{e} := \mathbf{e}/\varepsilon$, so that \mathbf{e} is an exponentially distributed r.v. with parameter ε which is independent of X. Then according to the previous equality,

$$\begin{split} \mathbb{E}_{x}(e^{-\lambda X_{\mathbf{e}}}\mathbb{I}_{\{\tau=\infty\}}) &= e^{-\lambda x}\mathbb{E}(e^{-\lambda X_{\mathbf{e}}}\mathbb{I}_{\{\underline{X}_{\mathbf{e}}\geq -x\}}(1-e^{-\rho(X_{\mathbf{e}}+x)})) \\ &= e^{-\lambda x}\mathbb{E}(e^{-\lambda X_{\mathbf{e}}}\mathbb{I}_{\{\underline{X}_{\mathbf{e}}\geq -x\}}) - e^{-(\lambda+\rho)x}\mathbb{E}(e^{-(\lambda+\rho)X_{\mathbf{e}}}\mathbb{I}_{\{\underline{X}_{\mathbf{e}}\geq -x\}}) \\ &= e^{-\lambda x}\mathbb{E}(e^{-\lambda(X_{\mathbf{e}}-\underline{X}_{\mathbf{e}}}))\mathbb{E}(e^{-\lambda\underline{X}_{\mathbf{e}}}\mathbb{I}_{\{\underline{X}_{\mathbf{e}}\geq -x\}}) - \\ &e^{-(\lambda+\rho)x}\mathbb{E}(e^{-(\lambda+\rho)(X_{\mathbf{e}}-\underline{X}_{\mathbf{e}})})\mathbb{E}(e^{-(\lambda+\rho)\underline{X}_{\mathbf{e}}}\mathbb{I}_{\{\underline{X}_{\mathbf{e}}\geq -x\}}), \end{split}$$

where the third equality above comes from the fact that \underline{X}_{e} and $X_{e} - \underline{X}_{e}$ are independent, see Th. 5, chap. VI in [1]. On the other hand recall that $-\underline{X}_{e}$ is exponentially distributed with parameter $\psi(\varepsilon)$ and that the law of $X_{e} - \underline{X}_{e}$ is given by

$$\mathbb{E}(e^{-\alpha(X_{\mathbf{e}}-\underline{X}_{\mathbf{e}})}) = \frac{\varepsilon}{\varepsilon - \varphi(\alpha)} \left(1 - \frac{\alpha}{\psi(\varepsilon)}\right),\,$$

see Th. 4, chap. VII in [1]. We derive from above that,

$$\mathbb{E}_{x}(e^{-\lambda X_{\mathbf{e}}}\mathbb{I}_{\{\tau=\infty\}}) = e^{-\lambda x} \frac{\varepsilon}{\varphi(\lambda) - \varepsilon} (e^{(\lambda - \psi(\varepsilon))x} - 1) -$$

$$e^{-(\lambda + \rho)x} \frac{\varepsilon}{\varphi(\lambda + \rho) - \varepsilon} (e^{((\lambda + \rho) - \psi(\varepsilon))x} - 1) .$$
(4.22)

On the other hand, note that

$$\begin{split} \mathbb{E}_x \left(\int_0^{\mathbf{e}} \frac{du}{X_u} \mathbb{I}_{\{\tau=\infty\}} \right) &= \mathbb{E}_x \left(\int_0^{\infty} \varepsilon e^{-\varepsilon y} dy \int_0^y \frac{du}{X_u} \mathbb{I}_{\{\tau=\infty\}} \right) \\ &= \mathbb{E}_x \left(\varepsilon \int_0^{\infty} \frac{du}{X_u} \int_u^{\infty} e^{-\varepsilon y} dy \mathbb{I}_{\{\tau=\infty\}} \right) \\ &= \mathbb{E}_x \left(\int_0^{\infty} \frac{e^{-\varepsilon u}}{X_u} du \mathbb{I}_{\{\tau=\infty\}} \right) = \mathbb{E}_x \left(\frac{1}{\varepsilon X_{\mathbf{e}}} \mathbb{I}_{\{\tau=\infty\}} \right) \,. \end{split}$$

Applying this remark and taking $\varepsilon = h(n)$ in the relation (4.22) once integrated over $(0, \infty)$ with respect to λ , yields

$$\mathbb{E}_x\left(\int_0^{\mathbf{e}_n} \frac{du}{X_u} \mathbb{1}_{\{\tau=\infty\}}\right) = \int_0^\infty \frac{e^{-\psi(h(n))x} - e^{-\lambda x}}{\varphi(\lambda) - h(n)} - \frac{e^{-\psi(h(n))x} - e^{-(\lambda+\rho)x}}{\varphi(\lambda+\rho) - h(n)} d\lambda, \quad (4.23)$$

where e_n is defined in the statement. We can check that the function $f(\lambda) := \frac{e^{-\psi(h(n))x} - e^{-\lambda x}}{\varphi(\lambda) - h(n)}$ is integrable over (0, M), for all M > 0 so that

$$\int_0^\infty f(\lambda) - f(\lambda + \rho) d\lambda = \lim_{M \to \infty} \int_0^M f(\lambda) d\lambda - \int_0^M f(\lambda + \rho) d\lambda$$
$$= \lim_{M \to \infty} \left(\int_0^\rho f(\lambda) d\lambda - \int_M^{M + \rho} f(\lambda) d\lambda \right) = \int_0^\rho f(\lambda) d\lambda \,.$$

This gives (4.20) and applying the same arguments to $X^{(n)}$ gives (4.21). Lemma 4.9. Let e_n be as in Lemma 4.8. Then for all x > 0,

$$\lim_{n \to \infty} \mathbb{E}_x \left(\int_0^{\mathbf{e}_n} \frac{du}{X_u} \mathbb{1}_{\{\tau = \infty\}} \right) = \lim_{n \to \infty} \mathbb{E}_x \left(\int_0^{\mathbf{e}_n} \frac{du}{X_u^{(n)}} \mathbb{1}_{\{\tau^{(n)} = \infty\}} \right)$$
$$= \mathbb{E}_x \left(\int_0^\infty \frac{du}{X_u} \mathbb{1}_{\{\tau = \infty\}} \right) = \int_0^\rho \frac{e^{-\lambda x} - e^{-\rho x}}{\phi(\lambda)} < \infty.$$
(4.24)

Proof. Fix x > 0, then the equality $\lim_{n\to\infty} \mathbb{E}_x \left(\int_0^{e_n} \frac{du}{X_u} \mathbb{1}_{\{\tau=\infty\}} \right) = \mathbb{E}_x \left(\int_0^\infty \frac{du}{X_u} \mathbb{1}_{\{\tau=\infty\}} \right)$ simply follows from monotone convergence. For the other equalities, thanks to Lemma 4.8 it suffices to show that,

$$\lim_{n \to \infty} \int_0^{\rho} \frac{e^{-\lambda x} - e^{-\psi(h(n))x}}{h(n) + \phi(\lambda)} d\lambda = \lim_{n \to \infty} \int_0^{\rho_n} \frac{e^{-\lambda x} - e^{-\psi^{(n)}(h(n))x}}{h(n) + \phi^{(n)}(\lambda)} d\lambda$$
$$= \int_0^{\rho} \frac{e^{-\lambda x} - e^{-\rho x}}{\phi(\lambda)} d\lambda < \infty.$$
(4.25)

We shall only prove that $\lim_{n\to\infty} \int_0^{\rho_n} \frac{e^{-\lambda x} - e^{-\psi^{(n)}(h(n))x}}{h(n) + \phi^{(n)}(\lambda)} d\lambda = \int_0^{\rho} \frac{e^{-\lambda x} - e^{-\rho x}}{\phi(\lambda)} d\lambda < \infty$. The first limit follows the same lines and is actually easier to obtain. Recall that for all n, $\rho_n \leq \psi^{(n)}(h(n))$, $\rho_n \leq \rho$ and that both sequences $(\psi^{(n)}(h(n)))$ and (ρ_n) converge to ρ . Then write,

$$\int_{0}^{\rho_{n}} \frac{e^{-\lambda x} - e^{-\psi^{(n)}(h(n))x}}{h(n) + \phi^{(n)}(\lambda)} d\lambda = \int_{0}^{\rho_{n}} \frac{e^{-\lambda x} - e^{-\rho_{n}x}}{h(n) + \phi^{(n)}(\lambda)} d\lambda$$

$$+ (e^{-\rho_{n}x} - e^{-\psi^{(n)}(h(n))x}) \int_{0}^{\rho_{n}} \frac{d\lambda}{h(n) + \phi^{(n)}(\lambda)}.$$
(4.26)

Note that $\phi^{(n)}(\rho_n) = 0$ and hence from the mean value theorem, for all $n \ge 0$ and $\lambda \in (0, \rho_n)$, there is $\alpha_n \in (\lambda, \rho_n)$ such that $\phi^{(n)}(\lambda) = (\rho_n - \lambda)|\phi^{(n)'}(\alpha_n)|$. But recall that (ρ_n) is increasing and that $\lim_n \rho_n = \rho$. Since $\lim_{n\to\infty} \phi^{(n)'} = \phi'$ and $\phi'(\rho) \in (0, \infty)$, there are c > 0, $\alpha \in (0, \rho_0)$ and $n_0 \ge 0$, such that for all $n \ge n_0$ and $\lambda \in (\rho_n - \alpha, \rho_n)$, $|\phi^{(n)'}(\lambda)| \ge c$. This implies that for all $n \ge n_0$ and all $\lambda \in (\rho_n - \alpha, \rho_n)$,

$$\frac{\rho_n - \lambda}{\phi^{(n)}(\lambda)} \le c^{-1} \,, \tag{4.27}$$

so that,

$$\int_{\rho_n-\alpha}^{\rho_n} \frac{d\lambda}{h(n) + \phi^{(n)}(\lambda)} = O(\log(h(n))).$$
(4.28)

Moreover, for all $\delta \in (0, 1)$ and n sufficiently large,

$$\int_{0}^{\rho_{n}-\alpha} \frac{d\lambda}{h(n)+\phi^{(n)}(\lambda)} = \int_{0}^{\delta h(n)} \frac{d\lambda}{h(n)+\phi^{(n)}(\lambda)} + \int_{\delta h(n)}^{\rho_{n}-\alpha} \frac{d\lambda}{\phi^{(n)}(\lambda)}$$
$$\leq (\operatorname{cst})\delta + \int_{\delta h(n)}^{\rho_{n}-\alpha} \frac{d\lambda}{\phi^{(n)}(\lambda)},$$

and $\int_{\delta h(n)}^{\rho_n - \alpha} \frac{d\lambda}{\phi^{(n)}(\lambda)}$ tends to $\int_0^{\rho - \alpha} \frac{d\lambda}{\phi(\lambda)}$, since $h \in \mathbb{Z}_0$. On the other hand, from Fatou's lemma,

$$\int_0^{\rho-\alpha} \frac{d\lambda}{\phi(\lambda)} \le \liminf_{n \to \infty} \int_0^{\rho_n-\alpha} \frac{d\lambda}{h(n) + \phi^{(n)}(\lambda)}$$

Since δ is chosen arbitrarily, we derive that

$$\lim_{n \to \infty} \int_0^{\rho_n - \alpha} \frac{d\lambda}{h(n) + \phi^{(n)}(\lambda)} = \int_0^{\rho - \alpha} \frac{d\lambda}{\phi(\lambda)} < \infty.$$
(4.29)

Then (4.29) together with (4.28) show that $\int_0^{\rho_n} \frac{d\lambda}{h(n) + \phi^{(n)}(\lambda)} = O(\log(h(n)))$. Moreover, since $\psi^{(n)}(0) = \rho_n$ and $\psi^{(n)'}(h(n)) \to \psi'(0) \in (0,\infty), \ \psi^{(n)}(h(n)) \sim \rho + h(n)\psi'(0)$, as

 $n \uparrow \infty$ and therefore, $e^{-\rho_n x} - e^{-\psi^{(n)}(h(n))x} \sim \psi'(0)h(n)$ as $n \to \infty$. Then we can conclude that the second term of the right hand side of (4.26) satisfies, $\lim_{n\to\infty} (e^{-\rho_n x} - e^{-\psi^{(n)}(h(n))x}) \int_0^{\rho_n} \frac{d\lambda}{h(n) + \phi^{(n)}(\lambda)} = 0.$

Let us now write the first term of the right hand side of (4.26) as

$$\int_0^{\rho_n} \frac{e^{-\lambda x} - e^{-\rho_n x}}{h(n) + \phi^{(n)}(\lambda)} d\lambda = \int_0^{\rho_n - \alpha} \frac{e^{-\lambda x} - e^{-\rho_n x}}{h(n) + \phi^{(n)}(\lambda)} d\lambda + \int_{\rho_n - \alpha}^{\rho_n} \frac{e^{-\lambda x} - e^{-\rho_n x}}{h(n) + \phi^{(n)}(\lambda)} d\lambda,$$

where α is as above and n is sufficiently large so that $\rho_n - \alpha > 0$. Then we prove that

$$\lim_{n \to \infty} \int_0^{\rho_n - \alpha} \frac{e^{-\lambda x} - e^{-\rho_n x}}{h(n) + \phi^{(n)}(\lambda)} d\lambda = \int_0^{\rho - \alpha} \frac{e^{-\lambda x} - e^{-\rho x}}{\phi(\lambda)} d\lambda$$

in the same way as for (4.29). For the second term, by using (4.27), we obtain that $\frac{e^{-\lambda x}-e^{-\rho_n x}}{h(n)+\phi^{(n)}(\lambda)} \mathbb{I}_{[\rho_n-\alpha,\rho_n]}(\lambda)$ is bounded by a constant and we derive from dominated convergence that

$$\lim_{n \to \infty} \int_{\rho_n - \alpha}^{\rho_n} \frac{e^{-\lambda x} - e^{-\rho_n x}}{h(n) + \phi^{(n)}(\lambda)} d\lambda = \int_{\rho - \alpha}^{\rho} \frac{e^{-\lambda x} - e^{-\rho x}}{\phi(\lambda)} d\lambda$$

Then we have proved that $\lim_{n\to\infty} \int_0^{\rho_n} \frac{e^{-\lambda x} - e^{-\psi^{(n)}(h(n))x}}{h(n) + \phi^{(n)}(\lambda)} d\lambda = \int_0^{\rho} \frac{e^{-\lambda x} - e^{-\rho x}}{\phi(\lambda)} d\lambda$. The fact that $\int_0^{\rho} \frac{e^{-\lambda x} - e^{-\rho x}}{\phi(\lambda)} d\lambda < \infty$ is a straightforward consequence of the behaviour of ϕ around ρ , that is, $1/\phi(\lambda) = O(1/(\rho - \lambda))$, as $\lambda \uparrow \rho$.

Corollary 4.1. Let e_n be as in Lemma 4.8 and recall that $h \in \mathbb{Z}_0$. Then for all x > 0,

$$\int_0^{\mathbf{e}_n} \frac{du}{X_u^{(n)}} \mathbb{1}_{\{\tau^{(n)}=\infty\}} \xrightarrow[n \to \infty]{} \zeta \mathbb{1}_{\{\zeta < \infty\}}.$$

Proof. Let us first observe that from the representation (2.6),

$$\zeta \mathbb{I}_{\{\zeta < \infty\}} = \int_0^\infty \frac{du}{X_u} \mathbb{I}_{\{\tau = \infty\}}, \qquad (4.30)$$

and write,

$$\begin{split} \left| \int_{0}^{\mathbf{e}_{n}} \frac{du}{X_{u}^{(n)}} \mathbb{1}_{\{\tau^{(n)}=\infty\}} - \zeta \mathbb{1}_{\{\zeta<\infty\}} \right| &\leq \left| \int_{0}^{\mathbf{e}_{n}} \frac{du}{X_{u}^{(n)}} \mathbb{1}_{\{\tau^{(n)}=\infty\}} - \int_{0}^{\mathbf{e}_{n}} \frac{du}{X_{u}} \mathbb{1}_{\{\tau=\infty\}} \right| + \int_{\mathbf{e}_{n}}^{\infty} \frac{du}{X_{u}} \mathbb{1}_{\{\tau=\infty\}} \\ &\leq \left(\int_{0}^{\mathbf{e}_{n}} \frac{du}{X_{u}^{(n)}} \mathbb{1}_{\{\tau^{(n)}=\infty\}} - \int_{0}^{\mathbf{e}_{n}} \frac{du}{X_{u}} \mathbb{1}_{\{\tau^{(n)}=\infty\}} \right) + \left(\int_{0}^{\mathbf{e}_{n}} \frac{du}{X_{u}} \mathbb{1}_{\{\tau=\infty\}} - \int_{0}^{\mathbf{e}_{n}} \frac{du}{X_{u}} \mathbb{1}_{\{\tau^{(n)}=\infty\}} \right) \\ &+ \int_{\mathbf{e}_{n}}^{\infty} \frac{du}{X_{u}} \mathbb{1}_{\{\tau=\infty\}}. \end{split}$$

From dominated convergence and Lemma 4.9, the last term of the right hand side of this inequality tends to 0 in expectation. (Note that since $X^{(n)} \leq X$ both terms between parentheses are non negative.) Then the expectation of each of the four terms between the parenthesis tends to $\mathbb{E}_x\left(\int_0^\infty \frac{du}{X_u}\mathbb{I}_{\{\tau=\infty\}}\right)$, as $n \to \infty$. For the first and the third term, this is a direct consequence of Lemma 4.9. It only remains to check that $\lim_{n\to\infty} \mathbb{E}_x\left(\int_0^{e_n} \frac{du}{X_u}\mathbb{I}_{\{\tau=\infty\}}\right) = \mathbb{E}_x\left(\int_0^\infty \frac{du}{X_u}\mathbb{I}_{\{\tau=\infty\}}\right)$. But again it follows from Lemma 4.9 and dominated convergence.

Let us recall some facts on scale functions that will be useful for the proof of the next lemma. We refer to [11] for more details. The scale function $W^{(n)}$ of $X^{(n)}$ is a continuous increasing function whose Laplace transform is given by

$$\int_{0}^{\infty} e^{-\lambda x} W^{(n)}(x) \, dx = \frac{1}{|\phi^{(n)}|(\lambda)} \,, \quad \lambda \ge \rho_n \,. \tag{4.31}$$

Recall also that this function admits the following representation,

$$W^{(n)}(x) = e^{\rho_n x} \int_0^x e^{-\rho_n u} U_n(du), \quad x \ge 0,$$
(4.32)

where U_n is the potential measure of the upward ladder height process $H^{(n)}$ of $X^{(n)}$, see (40) in [11]. Note also that we from a general result for subordinators, see Proposition III.1 in [1] and inequality (5) in its proof, there is a universal constant C such that,

$$U_n([0,x]) \le C \frac{|1/x - \rho_n|}{|\phi^{(n)}|(1/x)}, \quad x > 0.$$
(4.33)

In particular, C depends neither on x nor on n.

Lemma 4.10. There is a constant $C \in (0, \infty)$ such that for all $x \ge 0$ and $n \ge 1$,

$$\mathbb{E}_x\left(\int_0^\infty \frac{du}{X_u^{(n)}} \mathbb{1}_{\left\{\tau^{(n)}=\infty, X_u^{(n)}<1/h(n)\right\}}\right) \leq C.$$

If (x_n) is any sequence such that $x_n \ge 1/h(n)$ for all n, then

$$\lim_{n \to \infty} \mathbb{E}_{x_n} \left(\int_0^\infty \frac{du}{X_u^{(n)}} \mathbb{1}_{\left\{ \tau^{(n)} = \infty, X_u^{(n)} < 1/h(n) \right\}} \right) = 0.$$

Proof. Dividing the expression (4.22) for $X^{(n)}$ by ε and letting ε going to 0, gives, for all $\lambda \ge \rho_n$,

$$\mathbb{E}_{x}\left(\int_{0}^{\infty} e^{-\lambda X_{u}^{(n)}} \mathbb{1}_{\{\tau^{(n)}=\infty\}} du\right) = \frac{e^{-\rho_{n}x} - e^{-\lambda x}}{\varphi^{(n)}(\lambda)} - \frac{e^{-\rho_{n}x} - e^{-(\lambda+\rho_{n})x}}{\varphi^{(n)}(\lambda+\rho_{n})}$$
$$= \int_{0}^{\infty} e^{-\lambda y} (1 - e^{-\rho_{n}y}) \left(e^{-\rho_{n}x} W^{(n)}(y) - W^{(n)}(y-x) \mathbb{1}_{\{y \ge x\}}\right) dy, \tag{4.34}$$

where the second equality follows from (4.31) (for $X^{(n)}$). The set of functions $y \mapsto e^{-\lambda y}$, $\lambda \geq \rho_n$, for any fixed $n \geq 0$, is total in the vector space of continuous functions on $(0,\infty)$. Therefore, through classical arguments, we can extend the last identity to any non negative, measurable function defined on $(0,\infty)$. In particular, replacing $y \mapsto e^{-\lambda y}$ by $y \mapsto \frac{1}{y} \mathbb{1}_{\{y < 1/h(n)\}}$ gives,

$$\mathbb{E}_{x}\left(\int_{0}^{\infty} \frac{du}{X_{u}^{(n)}} \mathbb{1}_{\left\{\tau^{(n)}=\infty, X_{u}^{(n)}<1/h(n)\right\}}\right)$$

$$=\int_{0}^{1/h(n)} \frac{1-e^{-\rho_{n}y}}{y} \left(e^{-\rho_{n}x}W^{(n)}(y) - W^{(n)}(y-x)\mathbb{1}_{\left\{y\geq x\right\}}\right) dy$$
(4.35)

$$= \int_0^x \frac{1 - e^{-\rho_n y}}{y} e^{-\rho_n x} W^{(n)}(y) \, dy + \int_x^{1/h(n)} \frac{1 - e^{-\rho_n y}}{y} \left(e^{-\rho_n x} W^{(n)}(y) - W^{(n)}(y - x) \right) \, dy + \int_x^{1/h(n)} \frac{1 - e^{-\rho_n y}}{y} \left(e^{-\rho_n x} W^{(n)}(y) - W^{(n)}(y - x) \right) \, dy + \int_x^{1/h(n)} \frac{1 - e^{-\rho_n y}}{y} \left(e^{-\rho_n x} W^{(n)}(y) - W^{(n)}(y - x) \right) \, dy + \int_x^{1/h(n)} \frac{1 - e^{-\rho_n y}}{y} \left(e^{-\rho_n x} W^{(n)}(y) - W^{(n)}(y - x) \right) \, dy + \int_x^{1/h(n)} \frac{1 - e^{-\rho_n y}}{y} \left(e^{-\rho_n x} W^{(n)}(y) - W^{(n)}(y - x) \right) \, dy + \int_x^{1/h(n)} \frac{1 - e^{-\rho_n y}}{y} \left(e^{-\rho_n x} W^{(n)}(y) - W^{(n)}(y - x) \right) \, dy + \int_x^{1/h(n)} \frac{1 - e^{-\rho_n y}}{y} \left(e^{-\rho_n x} W^{(n)}(y) - W^{(n)}(y - x) \right) \, dy + \int_x^{1/h(n)} \frac{1 - e^{-\rho_n y}}{y} \left(e^{-\rho_n x} W^{(n)}(y) - W^{(n)}(y - x) \right) \, dy + \int_x^{1/h(n)} \frac{1 - e^{-\rho_n y}}{y} \left(e^{-\rho_n x} W^{(n)}(y) - W^{(n)}(y - x) \right) \, dy + \int_x^{1/h(n)} \frac{1 - e^{-\rho_n y}}{y} \left(e^{-\rho_n x} W^{(n)}(y) - W^{(n)}(y - x) \right) \, dy + \int_x^{1/h(n)} \frac{1 - e^{-\rho_n y}}{y} \left(e^{-\rho_n x} W^{(n)}(y) - W^{(n)}(y - x) \right) \, dy + \int_x^{1/h(n)} \frac{1 - e^{-\rho_n y}}{y} \left(e^{-\rho_n x} W^{(n)}(y) - W^{(n)}(y - x) \right) \, dy + \int_x^{1/h(n)} \frac{1 - e^{-\rho_n y}}{y} \left(e^{-\rho_n x} W^{(n)}(y) - W^{(n)}(y - x) \right) \, dy + \int_x^{1/h(n)} \frac{1 - e^{-\rho_n y}}{y} \left(e^{-\rho_n x} W^{(n)}(y) - W^{(n)}(y) \right) \, dy + \int_x^{1/h(n)} \frac{1 - e^{-\rho_n y}}{y} \left(e^{-\rho_n x} W^{(n)}(y) - W^{(n)}(y) \right) \, dy + \int_x^{1/h(n)} \frac{1 - e^{-\rho_n y}}{y} \left(e^{-\rho_n x} W^{(n)}(y) - W^{(n)}(y) \right) \, dy + \int_x^{1/h(n)} \frac{1 - e^{-\rho_n y}}{y} \left(e^{-\rho_n x} W^{(n)}(y) - W^{(n)}(y) \right) \, dy + \int_x^{1/h(n)} \frac{1 - e^{-\rho_n y}}{y} \left(e^{-\rho_n x} W^{(n)}(y) - W^{(n)}(y) \right) \, dy + \int_x^{1/h(n)} \frac{1 - e^{-\rho_n y}}{y} \left(e^{-\rho_n x} W^{(n)}(y) - W^{(n)}(y) \right) \, dy + \int_x^{1/h(n)} \frac{1 - e^{-\rho_n y}}{y} \left(e^{-\rho_n x} W^{(n)}(y) - W^{(n)}(y) \right) \, dy + \int_x^{1/h(n)} \frac{1 - e^{-\rho_n y}}{y} \left(e^{-\rho_n x} W^{(n)}(y) \right) \, dy + \int_x^{1/h(n)} \frac{1 - e^{-\rho_n y}}{y} \left(e^{-\rho_n x} W^{(n)}(y) \right) \, dy + \int_x^{1/h(n)} \frac{1 - e^{-\rho_n y}}{y} \left(e^{-\rho_n x} W^{(n)}(y) \right) \, dy + \int_x^{1/h(n)} \frac{1 - e^{-\rho_n y}}{y} \left(e^{-\rho_n x} W^{(n)}(y) \right) \, dy + \int_x^{1/h(n)} \frac{1 -$$

where we assume first that $x \leq 1/h(n)$. Note that the sequence $(\phi^{(n)'}(\rho_n)) = (\phi'(\rho_n + \varepsilon_n))$ tends to $\phi'(\rho) < 0$. Hence it is bounded away from 0 so that from (53) in the proof of Lemma 3.3 in [11], there is a constant C that depends neither on x nor on n, such that $W^{(n)}(x) \leq Ce^{\rho_n x}$, for all $x \geq 0$. Since (ρ_n) is increasing and tends to ρ , we obtain the bound,

$$\int_0^x \frac{1 - e^{-\rho_n y}}{y} e^{-\rho_n x} W^{(n)}(y) \, dy \le C \int_0^x \frac{1 - e^{-\rho y}}{y} e^{-\rho_0 (x-y)} \, dy \,, \tag{4.36}$$

and we readily show that the right hand side of this inequality tends to 0 as x tends to ∞ .

Now let us consider the second term of the last member of (4.35) and apply (4.32) in order to obtain,

$$\int_{x}^{1/h(n)} \frac{1 - e^{-\rho_n y}}{y} \left(e^{-\rho_n x} W^{(n)}(y) - W^{(n)}(y - x) \right) dy$$

=
$$\int_{x}^{1/h(n)} \frac{1 - e^{-\rho_n y}}{y} e^{-\rho_n x} \left(\int_{y-x}^{y} e^{-\rho_n (u-y)} U_n(du) \right) dy.$$
(4.37)

Assume first that x is less than some constant c > 0, that is $x \le c$, and write

$$\int_{x}^{1/h(n)} \frac{1 - e^{-\rho_{n}y}}{y} e^{-\rho_{n}x} \left(\int_{y-x}^{y} e^{-\rho_{n}(u-y)} U_{n}(du) \right) dy \leq \int_{x}^{1/h(n)} \frac{1 - e^{-\rho_{n}y}}{y} \int_{y-x}^{y} U_{n}(du) dy$$

$$\leq \int_{0}^{1/h(n)} \int_{u}^{u+x} \frac{1 - e^{-\rho y}}{y} dy U_{n}(du)$$

$$= \int_{0}^{1} \int_{u}^{u+x} \frac{1 - e^{-\rho y}}{y} dy U_{n}(du) + \int_{1}^{1/h(n)} \int_{u}^{u+x} \frac{1 - e^{-\rho y}}{y} dy U_{n}(du). \quad (4.38)$$

The function $(x, u) \mapsto \int_{u}^{u+x} \frac{1-e^{-\rho y}}{y} dy$ is continuous on $[0, c] \times [0, 1]$ and from the convergence of $X^{(n)}$ toward X we derive that the measure $U_n(du)$ converges weakly toward U(du). Hence the first term of (4.38) is bounded uniformly in $x \in [0, c]$ and $n \ge 0$. On the other hand, an integration by part gives for the second term,

$$\int_{1}^{1/h(n)} \int_{u}^{u+x} \frac{1-e^{-\rho y}}{y} \, dy \, U_{n}(du) \leq \int_{1}^{1/h(n)} \frac{c}{u} U_{n}(du)$$
$$= \int_{1}^{1/h(n)} \frac{c}{u^{2}} U_{n}([0,u]) \, du + c \frac{U_{n}([0,1/h(n)])}{1/h(n)} - c U_{n}([0,1]) \, ,$$

and (4.33) yields after a change of variable,

$$\int_{1}^{1/h(n)} \frac{c}{u} U_n(du) \le C\left(\int_{h(n)}^{1} \frac{v - \rho_n}{\varphi^{(n)}(v)} \, dv + (h(n) - \rho_n) \, \frac{h(n)}{\varphi^{(n)}(h(n))}\right) \,. \tag{4.39}$$

Our assumption on h (i.e. $h \in \mathbb{Z}_0$) and the fact that $\lim_{n\to\infty} \frac{h(n)}{\varphi^{(n)}(h(n))} = 0$ shows that the term above is also uniformly bounded in $x \in [0, c]$ and $n \ge 0$.

Assume now that $x \ge c$ and note that the right hand side of (4.37) is less than

Then on the one hand,

$$\begin{split} e^{-\rho_n x} \int_0^{c/2} \int_{u \lor x}^{(u+x) \land 1/h(n)} \frac{e^{\rho_n(y-u)}}{y} \, dy \, U_n(du) &= e^{-\rho_n x} \int_0^{c/2} \int_{u \lor x-u}^{(u+x) \land 1/h(n)-u} \frac{e^{\rho_n v}}{u+v} \, dv \, U_n(du) \\ &\leq e^{-\rho_n x} \int_0^{c/2} \int_{x-u}^x \frac{e^{\rho_n v}}{u+v} \, dv \, U_n(du) \\ &\leq \int_0^{c/2} \int_{x-c/2}^x \frac{1}{u+v} \, dv \, U_n(du) \\ &\leq \frac{c}{2} \int_0^{c/2} \frac{1}{u+x-c/2} \, U_n(du) \le U_n([0,c/2]) \, . \end{split}$$

Since $U_n([0, c/2])$ converges toward U([0, c/2]) this last term is uniformly bounded in $x \ge c$ and $n \ge 0$. On the other hand,

$$e^{-\rho_n x} \int_{c/2}^{1/h(n)} \int_{u \lor x}^{(u+x)\land 1/h(n)} \frac{e^{\rho_n(y-u)}}{y} \, dy \, U_n(du) = e^{-\rho_n x} \int_{c/2}^{1/h(n)} \int_{u \lor x-u}^{(u+x)\land 1/h(n)-u} \frac{e^{\rho_n v}}{u+v} \, dv \, U_n(du)$$

$$\leq e^{-\rho_n x} \int_{c/2}^{1/h(n)} \int_0^x \frac{e^{\rho_n v}}{u+v} \, dv \, U_n(du) \leq e^{-\rho_n x} \int_{c/2}^{1/h(n)} \frac{e^{\rho_n x}-1}{\rho_n u} \, U_n(du) \leq \int_{c/2}^{1/h(n)} \frac{1}{\rho_n u} \, U_n(du) \, .$$

But this last term can be bounded uniformly in $x \ge c$ and $n \ge 0$ exactly as in (4.39). Then we proved that the term in (4.35) is bounded by some constant for all n and x such that $x \le 1/h(n)$.

Now for n and x such that $x \ge 1/h(n)$, the term in (4.35) becomes

$$\int_0^x \frac{1 - e^{-\rho_n y}}{y} e^{-\rho_n x} W^{(n)}(y) \, dy$$

But then the bound obtained in (4.36) is still valid and gives the result in this case. The second assertion of the lemma also follows from this argument.

We define the first passage time above the level $x \ge 0$ of $X^{(n)}$ by

$$\tau_x^{(n)} = \inf\{t : X_t^{(n)} \ge x\}$$

Lemma 4.11. Let e_n be as in Lemma 4.8. Then, for all x > 0,

$$\lim_{n \to \infty} \int_{\tau_{1/h(n)}^{(n)}}^{\mathbf{e}_n} \frac{du}{X_u^{(n)}} \mathbb{1}_{\{\tau^{(n)} = \infty\}} \xrightarrow[n \to \infty]{} \frac{\mathrm{L}^1(\mathbb{P}_x)}{n \to \infty} 0.$$

Proof. Let us first show that

$$\lim_{n \to \infty} \int_{\tau_{1/h(n)}^{(n)}}^{\mathbf{e}_n} \frac{du}{X_u^{(n)}} \mathbb{I}_{\left\{\tau^{(n)} = \infty, \tau_{1/h(n)}^{(n)} < \mathbf{e}_n\right\}} \xrightarrow[n \to \infty]{} \frac{\mathrm{L}^1(\mathbb{P}_x)}{n \to \infty} \mathbf{0}.$$
(4.40)

Note that

$$\int_{\tau_{1/h(n)}^{(n)}}^{\mathbf{e}_{n}} \frac{du}{X_{u}^{(n)}} \mathbb{I}_{\left\{\tau^{(n)}=\infty,\,\tau_{1/h(n)}^{(n)}<\mathbf{e}_{n}\right\}} \leq \int_{\tau_{1/h(n)}^{(n)}}^{\tau_{1/h(n)}^{(n)}+\mathbf{e}_{n}} \frac{du}{X_{u}^{(n)}} \mathbb{I}_{\left\{\tau^{(n)}=\infty\right\}}.$$

Therefore, from the strong Markov property, the inequality $1/h(n) \leq X^{(n)}(\tau_{1/h(n)}^{(n)})$, \mathbb{P}_x -a.s. and Lemma 4.8,

$$\mathbb{E}_{x}\left(\int_{\tau_{1/h(n)}^{(n)}}^{\mathbf{e}_{n}} \frac{du}{X_{u}^{(n)}} \mathbb{I}_{\left\{\tau^{(n)}=\infty,\,\tau_{1/h(n)}^{(n)}<\mathbf{e}_{n}\right\}}\right) \leq E_{1/h(n)}\left(\int_{0}^{\mathbf{e}_{n}} \frac{du}{X_{u}^{(n)}} \mathbb{I}_{\left\{\tau^{(n)}=\infty\right\}}\right) \\ = \int_{0}^{\rho_{n}} \frac{e^{-\lambda n} - e^{-\psi^{(n)}(h(n))n}}{h(n) + \phi^{(n)}(\lambda)} d\lambda$$

and this last term tends to 0 from Lemma 4.8 and dominated convergence. Hence (4.40) is proved.

It remains to show that

$$\lim_{n \to \infty} \int_{\mathbf{e}_n}^{\tau_{1/h(n)}^{(n)}} \frac{du}{X_u^{(n)}} \mathbb{1}_{\left\{\tau^{(n)} = \infty, \tau_{1/h(n)}^{(n)} > \mathbf{e}_n\right\}} \xrightarrow[n \to \infty]{} \overset{\mathrm{L}^1(\mathbb{P}_x)}{\xrightarrow[n \to \infty]{}} 0.$$
(4.41)

Let us note that

$$\mathbb{E}_{x}\left(\int_{\mathbf{e}_{n}}^{\tau_{1/h(n)}^{(n)}} \frac{du}{X_{u}^{(n)}} \mathbb{I}_{\left\{\tau^{(n)}=\infty,\,\tau_{1/h(n)}^{(n)}>\mathbf{e}_{n}\right\}}\right) \leq \mathbb{E}_{x}\left(\int_{\mathbf{e}_{n}}^{\infty} \frac{du}{X_{u}^{(n)}} \mathbb{I}_{\left\{\tau^{(n)}=\infty,\,X_{u}^{(n)}<1/h(n)\right\}}\right)$$

and

$$\begin{split} & \mathbb{E}_{x}\left(\int_{\mathbf{e}_{n}}^{\infty}\frac{du}{X_{u}^{(n)}}\mathbb{I}_{\left\{\tau^{(n)}=\infty, X_{u}^{(n)}<1/h(n)\right\}}\right) = \mathbb{E}_{x}\left(\mathbb{I}_{\left\{\underline{X}_{e_{n}}^{(n)}\geq0\right\}}\left(\int_{0}^{\infty}\frac{du}{X_{u}^{(n)}}\mathbb{I}_{\left\{\tau^{(n)}=\infty, X_{u}^{(n)}<1/h(n)\right\}}\right) \circ \theta_{e_{n}}\right) \\ & = \mathbb{E}_{x}\left(\mathbb{I}_{\left\{\underline{X}_{e_{n}}^{(n)}\geq0\right\}}\mathbb{E}_{X_{e_{n}}^{(n)}}\left(\int_{0}^{\infty}\frac{du}{X_{u}^{(n)}}\mathbb{I}_{\left\{\tau^{(n)}=\infty, X_{u}^{(n)}<1/h(n)\right\}}\right)\right). \end{split}$$

From the law of large numbers for Lévy processes, for all $m \ge 0$, $\lim_{n\to\infty} h(n)X_{e_n}^{(m)} = E(X_1^{(m)}) = \phi^{(m)'}(0) \le \lim_{n\to\infty} h(n)X_{e_n}^{(n)}$, a.s. Since $\lim_{m\to\infty} \phi^{(m)'}(0) = \infty$, we obtain that $\lim_{n\to\infty} h(n)X_{e_n}^{(n)} = \infty$, a.s. Then (4.41) follows from Lemma 4.10 and dominated convergence.

We recall the definition of the first passage time above the level $x \ge 0$ of $Z^{(n)}$:

$$\sigma_x^{(n)} = \inf\{t : Z_t^{(n)} \ge x\}$$

Proof of Theorem 3.3. Thanks to Corollary 4.1 and Lemma 4.10, and using the decomposition,

$$\sigma_{1/h(n)}^{(n)} \mathbb{1}_{\{\tau^{(n)}=\infty\}} = \int_0^{\mathbf{e}_n} \frac{du}{X_u^{(n)}} \mathbb{1}_{\{\tau^{(n)}=\infty\}} - \int_{\tau_{1/h(n)}^{(n)}}^{\mathbf{e}_n} \frac{du}{X_u^{(n)}} \mathbb{1}_{\{\tau^{(n)}=\infty\}}$$

we obtain $\sigma_{1/h(n)}^{(n)} \mathbb{1}_{\{\tau^{(n)}=\infty\}} \xrightarrow{L_1} \zeta \mathbb{1}_{\{\zeta<\infty\}}$. Now observe that $\sigma_{1/h(n)}^{(n)} \mathbb{1}_{\{\tau^{(n)}=\infty\}} \leq \sigma_{1/h(n)}^{(n)} \mathbb{1}_{\{\sigma_{1/h(n)}^{(n)}<\infty\}}$, \mathbb{P}_x -a.s. Then we shall conclude by proving that,

$$\lim_{n \to \infty} \mathbb{E}_x \left(\sigma_{1/h(n)}^{(n)} \mathbb{1}_{\{\sigma_{1/h(n)}^{(n)} < \infty\}} \right) = \lim_{n \to \infty} \mathbb{E}_x \left(\sigma_{1/h(n)}^{(n)} \mathbb{1}_{\{\tau^{(n)} = \infty\}} \right) .$$
(4.42)

On the one hand, an application of the Markov property allows us to define the process $X^{(n)}$ conditioned to stay positive as follows: for all $t \ge 0$ and $\Lambda \in \mathcal{F}_t^{(n)}$,

$$\mathbb{E}_x(\mathbb{I}_{\Lambda} \mid \tau^{(n)} = \infty) = \mathbb{E}_x \left(\frac{1 - e^{-\rho_n X_t^{(n)}}}{1 - e^{-\rho_n x}} \mathbb{I}_{\Lambda} \mathbb{I}_{\{t < \tau^{(n)}\}} \right)$$

where $(\mathcal{F}_t^{(n)})$ is the natural filtration generated by the process $X^{(n)}$. Extending this formula from time t to the stopping time $\tau_{1/h(n)}^{(n)}$ and replacing \mathbb{I}_{Λ} by the $\mathcal{F}(\tau_{1/h(n)}^{(n)})$ -measurable functional $\int_0^{\tau_{1/h(n)}^{(n)}} du/X_u^{(n)} = \sigma_{1/h(n)}^{(n)}$ in the above formula allows us to write,

$$\mathbb{E}_{x}\left(\sigma_{1/h(n)}^{(n)} \mid \tau^{(n)} = \infty\right) = \mathbb{E}_{x}\left(\frac{1 - e^{-\rho_{n}X^{(n)}(\tau_{1/h(n)}^{(n)})}}{1 - e^{-\rho_{n}x}}\sigma_{1/h(n)}^{(n)}\mathbb{I}_{\{\tau_{1/h(n)}^{(n)} < \tau^{(n)}\}}\right).$$

Then from this relation together with the equality $\{\sigma_{1/h(n)}^{(n)} < \infty\} = \{\tau_{1/h(n)}^{(n)} < \tau^{(n)}\}$ and the inequality $X_{\tau_{1/h(n)}^{(n)}}^{(n)} \ge 1/h(n)$, we obtain,

$$\mathbb{E}_{x}\left(\sigma_{1/h(n)}^{(n)}\mathbb{I}_{\{\sigma_{1/h(n)}^{(n)}<\infty\}}\right) = \mathbb{E}_{x}\left(\sigma_{1/h(n)}^{(n)}\mathbb{I}_{\{\tau_{1/h(n)}^{(n)}<\tau^{(n)}\}}\right) \\
\leq \mathbb{E}_{x}\left(\frac{1-e^{-\rho_{n}x}}{1-e^{-\rho_{n}/h(n)}}\frac{1-e^{-\rho_{n}X^{(n)}(\tau_{1/h(n)}^{(n)})}}{1-e^{-\rho_{n}x}}\sigma_{1/h(n)}^{(n)}\mathbb{I}_{\{\tau_{1/h(n)}^{(n)}<\tau^{(n)}\}}\right) \\
= \frac{1}{1-e^{-\rho_{n}/h(n)}}\mathbb{E}_{x}\left(\sigma_{1/h(n)}^{(n)}\mathbb{1}_{\{\tau^{(n)}=\infty\}}\right).$$
(4.43)

On the other hand, the relation

$$\{\sigma_{1/h(n)}^{(n)} < \infty\} = \{\tau_{1/h(n)}^{(n)} < \tau^{(n)}\} = \{\tau^{(n)} = \infty\} \cup \{\tau_{1/h(n)}^{(n)} < \tau^{(n)}, \tau^{(n)} < \infty\},\$$

yields,

$$\mathbb{E}_{x}\left(\sigma_{1/h(n)}^{(n)}\mathbb{1}_{\{\tau^{(n)}=\infty\}}\right) \leq \mathbb{E}_{x}\left(\sigma_{1/h(n)}^{(n)}\mathbb{1}_{\{\sigma_{1/h(n)}^{(n)}<\infty\}}\right).$$

This inequality together with (4.43) allow us to obtain (4.42) and concludes the proof. \Box

4.3.2 Almost sure convergence

Lemma 4.12. For all x > 0, there exists a constant C such that for all $n \ge 0$,

$$\mathbb{E}_{x}\left(\tau_{1/h(n)}^{(n)}\mathbb{1}_{\{\tau^{(n)}=\infty\}}\right) \leq \frac{C}{\phi^{(n)}(h(n))}$$

Proof. If X is a subordinator, then $\mathbb{E}_x(\tau_{1/h(n)}^{(n)}\mathbb{1}_{\{\tau^{(n)}=\infty\}}) = \mathbb{E}_x(\tau_{1/h(n)}^{(n)})$, and the result follows from Proposition III.1 in [1]. If X is not a subordinator, then write

$$\mathbb{E}_x\left(\tau_{1/h(n)}^{(n)}\mathbb{1}_{\{\tau^{(n)}=\infty\}}\right) \le \mathbb{E}_x\left(\int_0^\infty \mathbb{1}_{\{X_t^{(n)}\leq 1/h(n)\}}\mathbb{1}_{\{\tau^{(n)}=\infty\}}dt\right).$$

As already justified in the beginning of the proof of Lemma 4.10, formula (4.34) may be extended to the function $y \mapsto \mathbb{1}_{\{y \leq 1/h(n)\}}$, so that

$$\mathbb{E}_x\left(\int_0^\infty \mathbb{1}_{\{X_t^{(n)} \le 1/h(n)\}} \mathbb{1}_{\{\tau^{(n)} = \infty\}}\right) = \int_0^{1/h(n)} (1 - e^{-\rho_n y}) \left(e^{-\rho_n x} W^{(n)}(y) - W^{(n)}(y - x) \mathbb{1}_{\{y \ge x\}}\right) dy$$

With calculations similar to $\left(4.36\right)$ and $\left(4.37\right)$, we find that this last term is less or equal than

$$c_x + \int_0^{1/h(n)} \int_u^{u+x} (1 - e^{-\rho y}) \, dy \, U_n(du) \, \le c_x + x U_n([0, 1/h(n)]) \\ \le c_x + c_1 x \frac{h(n) - \rho_n}{\varphi^{(n)}(h(n))} \le \frac{C}{\phi^{(n)}(h(n))} \,,$$

where c_x and c_1 are constants which do not depend on n and where we used (4.33) for the last inequality.

Lemma 4.13. Let e_n be as in Lemma 4.8.

1. If
$$\sum_{n\geq 0} \frac{\phi(\varepsilon_n)}{\phi(\varepsilon_n+h(n))} < \infty$$
, then $\left((X_{\tau_{1/h(n)}^{(n)}} - X_{\tau_{1/h(n)}^{(n)}}^{(n)}) \mathbb{1}_{\{\tau_n=\infty\}} \right)_{n\geq 0}$ converges toward 0, \mathbb{P}_x -almost surely, for all $x > 0$.

2. If
$$\sum_{n\geq 0} \frac{\phi(\varepsilon_n)}{h(n)} < \infty$$
, then $\left((X_{e_n} - X_{e_n}^{(n)}) \mathbb{1}_{\{\tau_n = \infty\}} \right)_{n\geq 0}$ converges toward 0, \mathbb{P}_x -almost surely, for all $x > 0$.

Proof. Let e be an exponentially distributed random variable with parameter 1 which is independent of the processes X and $X^{(n)}$, $n \ge 0$. To simplify our calculations, let us denote by $(Y_t^{(n)}, t \ge 0)$ the process $(X_t - X_t^{(n)}, t \ge 0)$ and recall that from Theorem 2.3, this process is a subordinator with Laplace exponent $\varphi - \varphi^{(n)}$ and that it is independent of $X^{(n)}$. Then for all a > 0 and $n \ge 0$,

$$\mathbb{P}_{x}\left(Y_{\tau_{1/h(n)}^{(n)}}^{(n)}\mathbb{1}_{\{\tau^{(n)}=\infty\}} > ae\right) = \int_{0}^{\infty} e^{-t}\mathbb{P}_{x}\left(Y_{\tau_{1/h(n)}^{(n)}}^{(n)}\mathbb{1}_{\{\tau^{(n)}=\infty\}} > at\right)dt \qquad (4.44)$$

$$\geq \int_{0}^{1} e^{-t}\mathbb{P}_{x}\left(Y_{\tau_{1/h(n)}^{(n)}}^{(n)}\mathbb{1}_{\{\tau^{(n)}=\infty\}} > a\right)dt \geq (1-e^{-1})\mathbb{P}_{x}\left(Y_{\tau_{1/h(n)}^{(n)}}^{(n)}\mathbb{1}_{\{\tau^{(n)}=\infty\}} > a\right).$$

On the other hand conditioning by e and then by $(\tau_{1/h(n)}^{(n)}, \tau^{(n)})$ yields,

$$\mathbb{P}_{x}\left(Y_{\tau_{1/h(n)}^{(n)}}^{(n)}\mathbb{1}_{\{\tau^{(n)}=\infty\}} > ae\right) = 1 - \mathbb{E}_{x}\left(\exp\left(-\frac{1}{a}Y_{\tau_{1/h(n)}^{(n)}}^{(n)}\mathbb{1}_{\{\tau^{(n)}=\infty\}}\right)\right)$$
$$= 1 - \mathbb{E}_{x}\left(\exp\left(-\frac{1}{a}Y_{\tau_{1/h(n)}^{(n)}}^{(n)}\right)\mathbb{1}_{\{\tau^{(n)}=\infty\}} + \mathbb{1}_{\{\tau^{(n)}<\infty\}}\right)$$
$$= \mathbb{E}_{x}\left(\left(1 - e^{-(\varphi(1/a) - \varphi^{(n)}(1/a))\tau_{1/h(n)}^{(n)}}\right)\mathbb{1}_{\{\tau^{(n)}=\infty\}}\right)$$

and this last quantity is less or equal than $(\varphi(1/a) - \varphi^{(n)}(1/a))\mathbb{E}_x(\tau_{1/h(n)}^{(n)}\mathbb{1}_{\{\tau^{(n)}=\infty\}})$. Now note that $\sum_{n\geq 0} \frac{\phi(\varepsilon_n)}{\phi^{(n)}(h(n))} < \infty$ if and only if $\sum_{n\geq 0} \frac{\phi(\varepsilon_n)}{\phi(\varepsilon_n + h(n))} < \infty$. Then using the series expansion $\varphi(1/a) - \varphi^{(n)}(1/a) = \varphi(\varepsilon_n) + \varepsilon_n \varphi'(1/a) + o(\varepsilon_n)$, the fact that $\varepsilon_n = o(\varphi(\varepsilon_n))$, Lemma 4.12 and the hypothesis give $\sum_{n\geq 1} \mathbb{P}_x \left(Y_{\tau_{1/h(n)}}^{(n)} \mathbb{1}_{\{\tau^{(n)}=\infty\}} > ae \right) < \infty$. The result follows from (4.44) and Borel Cantelli's lemma.

The proof of the almost sure convergence of the sequence $\left((X_{e_n} - X_{e_n}^{(n)}) \mathbb{1}_{\{\tau_n = \infty\}} \right)_{n \ge 0}$ toward 0 is very similar, so it is omitted.

Proof of Theorem 3.4. Let $n \ge 0$, recall the expression (4.30) and write $\sigma_{1/h(n)}^{(n)} \mathbb{1}_{\{\tau^{(n)}=\infty\}} - \zeta \mathbb{1}_{\{\zeta < \infty\}}$ as

$$\begin{split} \int_{0}^{\tau_{1/h(n)}^{(n)}} \frac{du}{X_{u}^{(n)}} \mathbb{1}_{\{\tau^{(n)}=\infty\}} - \int_{0}^{\tau_{1/h(n)}^{(n)}} \frac{du}{X_{u}} \mathbb{1}_{\{\tau^{(n)}=\infty\}} + \int_{0}^{\tau_{1/h(n)}^{(n)}} \frac{du}{X_{u}} (\mathbb{1}_{\{\tau^{(n)}=\infty\}} - \mathbb{1}_{\{\tau=\infty\}}) \\ &- \int_{\tau_{1/h(n)}^{(n)}}^{\infty} \frac{du}{X_{u}} \mathbb{1}_{\{\tau=\infty\}} \,. \end{split}$$

The last two terms tend \mathbb{P}_x -a.s to 0 as n goes to ∞ , from dominated convergence. For the two first terms, recall that the process $(X_t - X_t^{(n)}, t \ge 0)$ is a subordinator, so that

$$\int_{0}^{\tau_{1/h(n)}^{(n)}} \left(\frac{1}{X_{u}^{(n)}} - \frac{1}{X_{u}}\right) \mathbb{1}_{\{\tau^{(n)} = \infty\}} du \leq \frac{X_{\tau_{1/h(n)}^{(n)}} - X_{1/h(n)}^{(n)}}{\inf_{u \in [0, \tau_{1/h(n)}^{(n)}]} X_{u}^{(n)}} \int_{0}^{\tau_{1/h(n)}^{(n)}} \frac{du}{X_{u}} \mathbb{1}_{\{\tau^{(n)} = \infty\}} du$$

Then the right hand side of this inequality tends \mathbb{P}_x -a.s. to 0 since on the one hand, \mathbb{P}_x -a.s.,

$$\lim_{n \to \infty} \int_0^{\tau_{1/h(n)}^{(n)}} \frac{du}{X_u} \mathbb{1}_{\{\tau^{(n)} = \infty\}} du = \int_0^\infty \frac{du}{X_u} \mathbb{1}_{\{\tau = \infty\}} du < \infty ,$$
$$\lim_{n \to \infty} \frac{\mathbb{1}_{\{\tau^{(n)} = \infty\}}}{\inf_{u \in [0, \tau_{1/h(n)}^{(n)}]} X_u^{(n)}} = \frac{\mathbb{1}_{\{\tau = \infty\}}}{\inf_{u \in [0, \infty)} X_u} < \infty ,$$

where we used dominated convergence in the first equality, and on the other hand, from Lemma 4.13, $\lim_{n\to\infty} \left(X_{\tau_{1/h(n)}^{(n)}} - X_{\tau_{1/h(n)}^{(n)}}^{(n)} \right) \mathbb{1}_{\{\tau^{(n)}=\infty\}} = 0$. Then we have proved that $\lim_{n} \sigma_{1/h(n)}^{(n)} \mathbb{1}_{\{\tau^{(n)}=\infty\}} = \zeta \mathbb{1}_{\{\zeta<\infty\}}, \mathbb{P}_x$ -a.s. Now using the equality $\sigma_{1/h(n)}^{(n)} \mathbb{1}_{\{\sigma_{1/h(n)}^{(n)}<\infty\}} = \sigma_{1/h(n)}^{(n)} \mathbb{1}_{\{\tau^{(n)}=\infty\}} - \sigma_{1/h(n)}^{(n)} \mathbb{1}_{\{\tau_{1/h(n)}^{(n)}<\tau^{(n)},\tau^{(n)}<\infty\}}$ together with the fact that $\lim_{n} \mathbb{1}_{\{\tau_{1/h(n)}^{(n)}<\tau^{(n)},\tau^{(n)}<\infty\}} = 0, \mathbb{P}_x$ -a.s. allows us to obtain the

convergence $\lim_{n} \sigma_{1/h(n)}^{(n)} \mathbb{1}_{\{\sigma_{1/h(n)}^{(n)} < \infty\}} = \zeta \mathbb{1}_{\{\zeta < \infty\}}, \mathbb{P}_x$ -a.s.

Proof of Proposition 3.6. Let us first write,

γ

$$\tilde{\zeta}^{(n)} \mathbb{I}_{\{\tilde{\zeta}^{(n)}<\infty\}} - \int_{0}^{\mathbf{e}_{n}} \frac{du}{X_{u}} \mathbb{I}_{\{\tau=\infty\}} = \int_{0}^{\mathbf{e}_{n}} \frac{du}{X_{u}^{(n)}} \mathbb{I}_{\{\mathbf{e}_{n}<\tau^{(n)}\}} - \int_{0}^{\mathbf{e}_{n}} \frac{du}{X_{u}} \mathbb{I}_{\{\tau=\infty\}}$$
$$= \int_{0}^{\mathbf{e}_{n}} du \left(\frac{1}{X_{u}^{(n)}} - \frac{1}{X_{u}}\right) \mathbb{I}_{\{\mathbf{e}_{n}<\tau^{(n)}\}} + \int_{0}^{\mathbf{e}_{n}} \frac{du}{X_{u}} \left(\mathbb{I}_{\{\mathbf{e}_{n}<\tau^{(n)}\}} - \mathbb{I}_{\{\tau=\infty\}}\right). \quad (4.45)$$

Then from (3.5), the expression (4.30) and monotone convergence, it suffices to prove that the above term tends almost surely to 0 as n tends to ∞ . The sequence $(\tau^{(n)})$ is non decreasing and tends almost surely to τ . Moreover, for P-almost every ω , there is n_0 such that for all $n \ge n_0$, $\mathbb{I}_{\{\tau^{(n)}=\infty\}}(\omega) = \mathbb{I}_{\{\tau=\infty\}}(\omega)$ and since (e_n) tends almost surely to ∞ , for \mathbb{P} -almost every ω , there is n_1 , such that for all $n \ge n_1$,

$$\mathbb{I}_{\{\mathbf{e}_n < \tau^{(n)}\}}(\omega) = \mathbb{I}_{\{\mathbf{e}_n < \tau^{(n)}\} \cap \{\tau = \infty\}}(\omega) = \mathbb{I}_{\{\tau = \infty\}}(\omega),$$

from which we derive that

$$\lim_{n} \int_{0}^{\mathbf{e}_{n}} \frac{du}{X_{u}} \left(\mathbb{1}_{\{\mathbf{e}_{n} < \tau^{(n)}\}} - \mathbb{1}_{\{\tau = \infty\}} \right) = 0, \quad \mathbb{P}\text{-a.s}$$

On the other hand, note the bound of the first term in (4.45),

$$\int_{0}^{\mathbf{e}_{n}} du \left(\frac{1}{X_{u}^{(n)}} - \frac{1}{X_{u}} \right) \mathbb{1}_{\{\mathbf{e}_{n} < \tau^{(n)}\}} \leq \frac{X_{\mathbf{e}_{n}} - X_{\mathbf{e}_{n}}^{(n)}}{\inf_{u \in [0, \mathbf{e}_{n}]} X_{u}^{(n)}} \int_{0}^{\mathbf{e}_{n}} \frac{du}{X_{u}} \mathbb{1}_{\{\tau^{(n)} = \infty\}} du, \quad \mathbb{P}\text{-a.s.}$$

Then we conclude from Lemma 4.13 in a similar way to the proof of Theorem 3.4.

Acknowledgement Both authors are grateful for financial support from the ANR project "Rawabranch" number ANR-23-CE40-0008.

References

- [1] BERTOIN, J. Lévy processes Cambridge University Press, 1996.
- [2] BINGHAM, N.H.; GOLDIE, C.M.; TEUGELS, J.L. Regular variation Encyclopedia Math. Appl., 27, Cambridge University Press, Cambridge, 1989.
- [3] CABALLERO, M.E.; PÉREZ GARMENDIA, J.L.; URIBE BRAVO, G. A Lampertitype representation of continuous-state branching processes with immigration Ann. Probab. 41 (2013), no. 3A, 1585–1627.
- [4] DONEY, R.A.; MALLER, R.A. Stability of the overshoot for Lévy processes Ann. Probab. 30 (2002), no. 1, 188–212.
- [5] GREY, D.R. Asymptotic behaviour of continuous time, continuous state-space branching processes. J. Appl. Probability, 11 (1974), 669–677.
- [6] KYPRIANOU, A.E. Fluctuations of Lévy processes with applications Universitext Springer, Heidelberg, 2014,
- [7] LI, B.; FOUCART, C.; ZHOU, X. Behaviors near explosion of nonlinear CSBPs with regularly varying mechanisms *Preprint*, arXiv:2404.06145.
- [8] LI, P.-S. A continuous-state polynomial branching process. Stochastic Process. Appl. 129 (2019), no.8, 2941–2967.
- [9] LI, B.; ZHOU, X. On the explosion of a class of continuous-state nonlinear branching processes *Electron. J. Probab.* 26 (2021), Paper No. 148.
- [10] LI, Z. Measure-valued branching Markov processes Probab. Theory Stoch. Model., 103, Springer, Berlin, 2022.
- [11] KUZNETSOV, A; KYPRIANOU, A.E.; RIVERO, V. The theory of scale functions for spectrally negative Lévy processes. Lecture Notes in Math. 2061, Lévy Matters, *Springer*, Heidelberg, 2012, 97–186.